

## Stable equivalence of Morita type and Frobenius extensions

MARGARET BEATTIE, STEFAAN CAENEPEEL AND ȘERBAN RAIANU

*Communicated by Constantin Năstăsescu*

**Abstract** - A.S. Dugas and R. Martínez-Villa proved that if there exists a stable equivalence of Morita type between the  $k$ -algebras  $\Lambda$  and  $\Gamma$ , then it is possible to replace  $\Lambda$  by a Morita equivalent  $k$ -algebra  $\Delta$  such that  $\Gamma$  is a subring of  $\Delta$  and the induction and restriction functors induce inverse stable equivalences. In this note we give an affirmative answer to a question of Alex Dugas about the existence of a  $\Gamma$ -coring structure on  $\Delta$ . We do this by showing that  $\Delta$  is a Frobenius extension of  $\Gamma$ .

**Key words and phrases** : stable equivalence of Morita type, Hopf algebra, Frobenius extension, coring.

**Mathematics Subject Classification** (2000) : primary 16D90; secondary 16D20, 16D50, 16W30, 16S40.

As in [4], we will assume throughout that the algebras  $\Lambda$  and  $\Gamma$  are finite dimensional over a field  $k$  and have no semisimple blocks.

The algebras  $\Lambda$  and  $\Gamma$  are said to be stably equivalent if the categories of finitely generated modules modulo projectives for  $\Lambda$  and  $\Gamma$  are equivalent (see [1]).

A pair of left-right projective bimodules  ${}_{\Lambda}M_{\Gamma}$  and  ${}_{\Gamma}N_{\Lambda}$  is said to induce a stable equivalence of Morita type between  $\Lambda$  and  $\Gamma$  if we have the following isomorphisms of bimodules:

$${}_{\Lambda}M \otimes_{\Gamma} N_{\Lambda} \simeq {}_{\Lambda}\Lambda_{\Lambda} \oplus {}_{\Lambda}P_{\Lambda} \quad \text{and} \quad {}_{\Gamma}N \otimes_{\Lambda} M_{\Gamma} \simeq {}_{\Gamma}\Gamma_{\Gamma} \oplus {}_{\Gamma}Q_{\Gamma}$$

where  ${}_{\Lambda}P_{\Lambda}$  and  ${}_{\Gamma}Q_{\Gamma}$  are projective bimodules (see [2]).

We begin by stating the result of Dugas and Martínez-Villa mentioned in the abstract:

**Theorem 1.1.** (see [4, Corollary 5.1]) *Let  $\Lambda$  and  $\Gamma$  be finite dimensional  $k$ -algebras whose semisimple quotients are separable. If at least one of them is indecomposable, then the following are equivalent:*

- (1) *There exists a stable equivalence of Morita type between  $\Lambda$  and  $\Gamma$ .*
- (2) *There exists a  $k$ -algebra  $\Delta$ , Morita equivalent to  $\Lambda$ , and an injective ring homomorphism  $\Gamma \hookrightarrow \Delta$  such that the restriction and induction functors are*

exact and induce inverse stable equivalences.

(3) There exists a  $k$ -algebra  $\Delta$ , Morita equivalent to  $\Lambda$ , and an injective ring homomorphism  $\Gamma \hookrightarrow \Delta$  such that

$${}_{\Gamma}\Delta_{\Gamma} = {}_{\Gamma}\Gamma_{\Gamma} \oplus {}_{\Gamma}P_{\Gamma} \quad \text{and} \quad {}_{\Delta}\Delta \otimes_{\Gamma} \Delta_{\Delta} \simeq {}_{\Delta}\Delta_{\Delta} \oplus {}_{\Delta}Q_{\Delta}$$

for projective bimodules  ${}_{\Gamma}P_{\Gamma}$  and  ${}_{\Delta}Q_{\Delta}$ .

We recall now the definition of Frobenius extension, and its dual notion, Frobenius coring.

**Definition 1.1.** (see [5]) *Let  $i : R \rightarrow S$  be a ring homomorphism. Then  $S/R$  is called a Frobenius extension if one of the following equivalent conditions is satisfied:*

1.  $S$  is finitely generated and projective as a right  $R$ -module and  $\text{Hom}_R(S, R)$  and  $S$  are isomorphic as  $(R, S)$ -bimodules.
2. There exists a Frobenius system  $(e, \varepsilon)$ , consisting of

$$e = e^1 \otimes e^2 \in (S \otimes_R S)^S = \{e^1 \otimes e^2 \in S \otimes_R S \mid se^1 \otimes e^2 = e^1 \otimes e^2 s, \forall s \in S\}$$

$$\text{and } \varepsilon : S \rightarrow R \text{ an } R\text{-bimodule map such that } \varepsilon(e^1)e^2 = e^1\varepsilon(e^2) = 1.$$

For the proof of the equivalence of the two conditions, see for example [3, Theorem 28].

**Definition 1.2.** (see [7]) *If  $R$  is a ring, a coring is a comonoid in the monoidal category of  $R$ -bimodules. So a coring consists of an  $R$ -bimodule  $C$ , together with a coassociative comultiplication  $C \rightarrow C \otimes_R C$  and counit  $C \rightarrow R$  which are both  $R$ -bimodule maps.*

$C$  is called a Frobenius  $R$ -coring if there exists a Frobenius system  $(\theta, 1)$ , consisting of an element  $1 \in C$  and an  $R$ -bimodule map  $\theta : C \otimes_R C \rightarrow R$  satisfying the conditions

$$c_{(1)}\theta(c_{(2)} \otimes d) = \theta(c \otimes d_{(1)})d_{(2)} \quad \text{and} \quad \theta(c \otimes 1) = \theta(1 \otimes c) = \varepsilon(c).$$

Let  $(S, m, 1, e, \varepsilon)$  be a Frobenius extension of  $R$ , and consider  $\Delta : S \rightarrow S \otimes_R S$ ,  $\Delta(s) = se = es$ . An easy verification shows that  $(S, \Delta, \varepsilon, \theta = \varepsilon \circ m, 1)$  is a Frobenius coring.

Conversely, if  $(C, \Delta, \varepsilon, \theta, 1)$  is a Frobenius  $R$ -coring, then  $(C, m, 1, \Delta(1), \varepsilon)$  is a Frobenius extension. Here  $m : C \otimes_R C \rightarrow C$ ,  $m(c \otimes d) = c_{(1)}\theta(c_{(2)} \otimes d) = \theta(c \otimes d_{(1)})d_{(2)}$ .

These two assertions basically tell us that Frobenius extension structures on an  $R$ -bimodule  $M$  correspond bijectively to Frobenius  $R$ -coring structures on  $M$ .

Let  $S$  be a Frobenius extension. Then the categories  $\mathcal{M}_S$  and  $\mathcal{M}^S$  are isomorphic: on a right  $S$ -module, we define a right  $S$ -coaction by  $\rho(m) = me^1 \otimes e^2$ . On a right  $S$ -comodule, we define a right  $S$ -action  $ms = m_{[0]}\varepsilon(m_{[1]}s)$ . The restriction functor  $G : \mathcal{M}_S \rightarrow \mathcal{M}_R$  has a left adjoint, the induction functor  $F$ ; the forgetful functor  $\mathcal{M}_S \rightarrow \mathcal{M}_R$  has a right adjoint. These functors are compatible with the above isomorphism. This implies that  $G$  is at the same time a left and a right adjoint of  $F$ .

**Definition 1.3.** (see [6] or [3, p.91]) *Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a covariant functor. If there exists a functor  $G : \mathcal{D} \rightarrow \mathcal{C}$  which is at the same time a right and a left adjoint of  $F$ , then we call  $F$  a Frobenius functor, and we say that  $(F, G)$  is a Frobenius pair for  $\mathcal{C}$  and  $\mathcal{D}$ .*

**Remark 1.1.** (see [5] or [3, Theorem 28, p.103]) *Let  $i : R \rightarrow S$  be a ring homomorphism,  $F$  the induction functor and  $G$  the restriction functor. If  $S/R$  is a Frobenius extension, then we have seen above that  $(F, G)$  is a Frobenius pair; in fact, it can be shown that the converse also holds:  $(F, G)$  is a Frobenius pair if and only if  $S/R$  is a Frobenius extension.*

We can now state and prove our result. Assertion (3) gives an affirmative answer to a question asked by Alex Dugas.

**Theorem 1.2.** *Let  $\Lambda$  and  $\Gamma$  be finite dimensional  $k$ -algebras whose semisimple quotients are separable. Assume that at least one of them is indecomposable, and that there exists a stable equivalence of Morita type between  $\Lambda$  and  $\Gamma$ . Then the following assertions hold:*

- (1) *There exists a  $k$ -algebra  $\Delta$ , Morita equivalent to  $\Lambda$ , and an injective ring homomorphism  $\Gamma \hookrightarrow \Delta$  such that the restriction and induction functors are a Frobenius pair.*
- (2) *There exists a  $k$ -algebra  $\Delta$ , Morita equivalent to  $\Lambda$ , and an injective ring homomorphism  $\Gamma \hookrightarrow \Delta$  such that  $\Delta/\Gamma$  is a Frobenius extension.*
- (3) *There exists a  $k$ -algebra  $\Delta$ , Morita equivalent to  $\Lambda$ , and an injective ring homomorphism  $\Gamma \hookrightarrow \Delta$  such that*

$${}_{\Gamma}\Delta_{\Gamma} = {}_{\Gamma}\Gamma_{\Gamma} \oplus {}_{\Gamma}P_{\Gamma} \quad \text{and} \quad {}_{\Delta}\Delta \otimes_{\Gamma} \Delta_{\Delta} \simeq {}_{\Delta}\Delta_{\Delta} \oplus {}_{\Delta}Q_{\Delta}$$

*for projective bimodules  ${}_{\Gamma}P_{\Gamma}$  and  ${}_{\Delta}Q_{\Delta}$ , and  $\Delta$  is a Frobenius  $\Gamma$ -coring with multiplication given by the injection of  ${}_{\Delta}\Delta_{\Delta}$  into  ${}_{\Delta}\Delta \otimes_{\Gamma} \Delta_{\Delta}$ , and counit given by the projection of  ${}_{\Gamma}\Delta_{\Gamma}$  onto  ${}_{\Gamma}\Gamma_{\Gamma}$ .*

**Proof.** (1) Suppose  ${}_{\Lambda}M_{\Gamma}$  and  ${}_{\Gamma}N_{\Delta}$  are indecomposable bimodules that induce a stable equivalence of Morita type. Let  $\Delta = \text{End}_{\Lambda}(M)$ . By the proof of (1) $\Rightarrow$ (2) of [4, Corollary 5.1], we have that

$$\text{Res}_{\Gamma}^{\Delta} \simeq (- \otimes_{\Lambda} M_{\Gamma}) \circ \text{Hom}_{\Delta}(M, -)$$

and

$$\text{Ind}_{\Gamma}^{\Delta} \simeq (- \otimes_{\Lambda} M_{\Delta}) \circ (- \otimes_{\Gamma} N_{\Lambda}).$$

Now  $- \otimes_{\Lambda} M_{\Gamma}$  is a right and left adjoint of  $- \otimes_{\Gamma} N_{\Lambda}$  by [4, Corollary 3.1,(2)], and  $\text{Hom}_{\Delta}(M, -)$  is a right and left adjoint of  $- \otimes_{\Lambda} M_{\Delta}$  because they are inverse equivalences, so  $\text{Res}_{\Gamma}^{\Delta}$  is a right and left adjoint of  $\text{Ind}_{\Gamma}^{\Delta}$ .

(2) follows from (1) and Remark 1.1.

(3) follows immediately from the above observation that a Frobenius extension is also a Frobenius coring.  $\square$

### Acknowledgements

The work on this note was started while the last two authors were visiting the Mathematics Department at Mount Allison University. They thank the department for its warm hospitality.

### References

- [1] M. AUSLANDER and I. REITEN, *Stable equivalences of artin algebras*, Proc. Conf. on Orders, Group Rings and Related Topics, Lecture Notes in Math., 353, Springer-Verlag, New York, 1973.
- [2] M. BROUÉ, *Equivalences of blocks of group algebras*, in: Finite Dimensional Algebras and related Topics, pp. 1-26, Kluwer, 1994.
- [3] S. CAENEPEEL, G. MILITARU and S. ZHU, *Frobenius and separable functors for generalized module categories and nonlinear equations* Lecture Notes in Math., 1787, Springer-Verlag, Berlin, 2002.
- [4] A.S. DUGAS and R. MARTÍNEZ-VILLA, A note on stable equivalences of Morita type, *J. Pure Appl. Algebra*, **208** (2007), 421-433.
- [5] F. KASCH, Projektive Frobenius-Erweiterungen, *S.-B. Heidelberger Akad. Wiss. (Math.-Nat. Kl.)*, **61** (1961), 89-109.
- [6] K. MORITA, Adjoint pairs of functors and Frobenius extensions, *Sci. Rep. Tokyo Kyoiku Daigaku (Sect. A)*, **9** (1965), 40-71.
- [7] M. SWEEDLER, The predual theorem to the Jacobson-Bourbaki theorem, *Trans. Amer. Math. Soc.*, **213** (1975), 391-406.

*Margaret Beattie*

Department of Mathematics and Computer Science, Mount Allison University  
Sackville, New Brunswick, Canada E4L 1E6  
E-mail: mbeattie@mta.ca

*Stefaan Caenepeel*

Faculteit Ingenieurswetenschappen, Vrije Universiteit Brussel  
Pleinlaan 2, B-1050 Brussel, Belgium  
E-mail: scaenep@vub.ac.be

*Șerban Raianu*

Mathematics Department, California State University  
Dominguez Hills, 1000 E Victoria St, Carson, CA 90747, USA  
E-mail: sraianu@csudh.edu