

PRE-PUBLICACIONES del seminario matemático

2008

“garcía de galdeano”

On complex infinite-dimensional
Grassmann manifolds

Daniel Beltita
José E. Galé

n. 10



seminario
matemático

garcía de galdeano

Universidad de Zaragoza

ON COMPLEX INFINITE-DIMENSIONAL GRASSMANN MANIFOLDS

DANIEL BELTIȚĂ AND JOSÉ E. GALÉ

ABSTRACT. We investigate geometric properties of Grassmann manifolds and their complexifications in an infinite-dimensional setting. Specific structures of quaternionic type are constructed on these complexifications by a direct method that does not require any use of the cotangent bundles.

1. INTRODUCTION

In this paper we discuss differential geometric properties of Grassmann manifolds in a C^* -algebraic setting. Background information for the circle of ideas centered on such manifolds here can be found in the works [PR87], [S88], [CPR90], [W90], [MR92], [MS98], [DG01], [DG02], [G06], and [B06], [U85]. Our approach to the subject originated in some specific questions concerning holomorphy, which in a natural way led us to investigate complex structures as well as complexifications of these manifolds.

Let \mathcal{H} be a *complex* Hilbert space. Recall that the Grassmannian $\text{Gr}(\mathcal{H})$ associated with \mathcal{H} , or, alternatively, with the Banach algebra $\mathcal{B}(\mathcal{H})$ of bounded operators on \mathcal{H} , is formed by all closed linear subspaces of \mathcal{H} and it is a complex Banach manifold (see [U85]). For more general C^* -algebras the definition of the Grassmann manifold is a bit more involved, and not always standard. For instance, if A is a unital C^* -algebra, the Grassmannian of A is sometimes assumed to be the set $\mathcal{P}_{sa}(A)$ of selfadjoint idempotents of A , endowed with the relative topology. This is so considered for example in [S88], where such Grassmannians are studied in connection with Hermitian holomorphic bundles as motivated by certain aspects of the Cowen-Douglas theory. The manifold $\mathcal{P}_{sa}(A)$ is in principle *real*-analytic, so that one must be careful in order to deal with holomorphy, see [S88], p. 278. In other settings it is suitable to deal with explicit complex Grassmann manifolds, as for example in [PR87] or [DG01], [DG02]. This is accomplished in the following way.

Recall that, for a given unital, complex, associative (C^* -) algebra A , two idempotents $p, q \in A$ are said to be equivalent if $pq = q$ and $qp = p$. The equivalence class of an idempotent p is denoted here by $[p]$, and the quotient set formed by all the classes $[p]$ is denoted by $\text{Gr}(A)$ and called the Grassmannian of A . Let G_A be the group of invertible elements of A and let U_A be the subgroup of G_A of unitary elements. Then, see Section 2 below, G_A has a natural action on $\text{Gr}(A)$. The isotropy subgroup of G_A at $[p]$ for such an action is denoted by $G_A([p])$. Then the G_A -orbits in $\text{Gr}(A)$ coincide with the quotients $G_A/G_A([p])$, which, endowed with their respective quotient topologies, are holomorphic Banach manifolds [DG01]. In analogy, put $G_A(p) := \{u^{-1}pu : u \in G_A\}$ and $U_A(p) := \{u^{-1}pu : u \in U_A\}$. As above, $G_A/G_A(p)$ is a complex Banach manifold and $U_A/U_A(p)$ is, for $p \in \mathcal{P}_{sa}(A)$, a real-analytic Banach manifold, in their respective quotient topologies. Let us remark that $U_A/U_A(p) = U_A/U_A([p])$ for every $p \in \mathcal{P}_{sa}(A)$ (see Remark 2.5) and that $\mathcal{P}_{sa}(A) \equiv \bigcup_{p \in \mathcal{P}_{sa}(A)} U_A/U_A([p])$. The complex structures of the Grassmannian $\text{Gr}(A) \equiv \bigcup G_A/U_G([p])$ and of its associated (holomorphic) Stiefel bundle have been plainly used in [DG01] and [DG02] in order to obtain holomorphic parametrizations of framings for projections on a fixed Banach space. It sounds sensible to analyze the relationship between the differentiable structures of the two types of Grassmannians $U_A/U_A(p)$ and $G_A/G_A([p])$ when $p = p^*$.

Another motivation in writing the present paper came from representation theory. To explain this, recall the classical setting of the Bott-Borel-Weil theorem of [Bo57] involving the flag manifolds and

Date: May 15, 2008.

2000 *Mathematics Subject Classification.* Primary 46L05; Secondary 58B12; 53C15; 43A85; 22E65.

Key words and phrases. Grassmann manifold; homogeneous vector bundle; complexification; hypercomplex structure; C^* -algebra.

realizations of representations on holomorphic sections spaces of holomorphic bundles. The notion of *complexification* plays a central role in this area, inasmuch as one of the ways to describe the complex structure of the flag manifolds is to view the latter as homogeneous spaces of complexifications of compact Lie groups.

A similar circle of ideas can be found in infinite dimensions for a class of manifolds which play a central role in many areas of functional analysis and operator theory. Specifically, let $\mathbf{1} \in B \subseteq A$ be unital C^* -algebras such that there exists a conditional expectation $E: A \rightarrow B$. Let φ be a state of A such that $\varphi \circ E = \varphi$, and let π_φ denote the Gelfand-Naimark-Segal (GNS, for short) representation obtained out of φ . Then it is possible to realize π_φ as acting on spaces of real-analytic sections of a certain homogeneous Hermitian vector bundle on U_A/U_B . Further, in some cases involving finiteness properties of spectra and traces of elements in A , one can prove that the homogeneous space U_A/U_B is a *complex* manifold as well and that the realization space is formed by *holomorphic* sections (see Theorem 5.4 and Theorem 5.8 in [BR07]).

Apart from the above result, the holomorphic character of the manifolds U_A/U_B and associated bundles is far from being clear in general. Since the Grassmannians $U_A/U_A(p)$ and tautological bundles over $U_A/U_A(p)$ are universal objects (see for example [ALRS97], [DG02] in connection with ideas here), we search for related results of holomorphy in this special case. Using a characterization (and labelling) of invariant complex structures on infinite-dimensional homogeneous manifolds (Theorem 6.1 in [B06]), we prove in Section 3 below that $U_A/U_A(p)$ and $G_A/G_A([p])$ are locally biholomorphic complex manifolds. Moreover, it is also shown that $G_A/G_A(p)$ is a complexification of $U_A/U_A(p)$, and then these results are translated in terms of homogeneous vector bundles, in the spirit of Theorem 5.8 in [BR07]. (A complementary perspective on these manifolds can be found in [BN05].) Section 4 is devoted to exhibit how the above results look like in the (on the other hand well known) case of the algebra $A = \mathcal{B}(\mathcal{H})$ and corresponding universal, tautological vector bundles.

Finally, $G_A/G_A(p)$ being a complexification of the complex manifold $U_A/U_A(p)$, we point out the existence of quaternionic structures for the above complexifications in Section 5. The occurrence of quaternionic structures on this level is a fairly known phenomenon in finite dimensions; see for instance the complexifications of Hermitian symmetric spaces of compact type studied in [BG96] and [BG98]. An infinite-dimensional version of this phenomenon was discussed in [T07] in the special case of the restricted Grassmann manifold. The latter manifold is modeled on a Hilbert space and is endowed with a Riemannian structure which allows one to construct almost complex structures on the tangent bundle by identifying it with the cotangent bundle.

Nothing of this kind is available in the case of the C^* -algebraic Grassmann manifolds investigated in the present paper. Instead, we have to construct the almost complex structures in a direct manner inspired by some of the earliest insights into the geometry of the tangent bundles; see [H60] and [Do62]. This approach leads to almost hypercomplex structures on the complexifications of the C^* -algebraic Grassmann manifolds provided by the tangent bundles and is related to the theory of adapted complex structures developed in finite dimensions in papers like [LS91], [Sz04], and [Bi04].

2. GRASSMANN MANIFOLDS IN AN ALGEBRAIC SETTING

We begin with several elementary considerations about idempotents in complex associative algebras.

Notation 2.1. We are going to use the following notation: A is a unital associative algebra over \mathbb{C} with unit $\mathbf{1}$ and set of idempotents $\mathcal{P}(A) = \{p \in A \mid p^2 = p\}$; for $p_1, p_2 \in \mathcal{P}(A)$ the notation $p_1 \sim p_2$ means that we have both $p_1 p_2 = p_2$ and $p_2 p_1 = p_1$. For each $p \in \mathcal{P}(A)$ we denote its equivalence class by $[p] := \{q \in \mathcal{P}(A) \mid q \sim p\}$. The quotient set is denoted by $\text{Gr}(A) = \mathcal{P}(A)/\sim$ (the *Grassmannian* of A) and the quotient map by $\pi: p \mapsto [p]$, $\mathcal{P}(A) \rightarrow \text{Gr}(A)$.

The group of invertible elements of A is denoted by G_A , and it has a natural action on $\mathcal{P}(A)$ by

$$\alpha: (u, q) \mapsto uqu^{-1}, \quad G_A \times \mathcal{P}(A) \rightarrow \mathcal{P}(A).$$

The corresponding isotropy group at $p \in \mathcal{P}(A)$ is $\{u \in G_A \mid \alpha(u, p) = p\} = G_A \cap \{p\}' = G_{\{p\}'} =: G(p)$ where we denote by $\{p\}'$ the commutant subalgebra of p in A (see page 484 in [DG02]). \square

Lemma 2.2. *There exists a well-defined action of the group G_A upon $\text{Gr}(A)$ like this:*

$$\beta: (u, [p]) \mapsto [upu^{-1}], \quad G_A \times \text{Gr}(A) \rightarrow \text{Gr}(A),$$

and the diagram

$$\begin{array}{ccc} G_A \times \mathcal{P}(A) & \xrightarrow{\alpha} & \mathcal{P}(A) \\ \text{id}_G \times \pi \downarrow & & \downarrow \pi \\ G_A \times \text{Gr}(A) & \xrightarrow{\beta} & \text{Gr}(A) \end{array}$$

is commutative.

Proof. See for instance the end of Section 3 in [DG01]. \square

Definition 2.3. For every idempotent $p \in \mathcal{P}(A)$ we denote by $G_A([p])$ the isotropy group of the action $\beta: G_A \times \text{Gr}(A) \rightarrow \text{Gr}(A)$ at the point $[p] \in \text{Gr}(A)$, that is, $G_A([p]) = \{u \in G_A \mid [upu^{-1}] = [p]\}$. \square

The following statement concerns the relationship between the isotropy groups of the actions α and β of G_A upon $\mathcal{P}(A)$ and $\text{Gr}(A)$, respectively.

Proposition 2.4. *The following assertions hold.*

- (i) *For every $p \in \mathcal{P}(A)$ we have $G_A([p]) \cap G_A([\mathbf{1} - p]) = G(p)$.*
- (ii) *If U is a subgroup of G_A and $p \in \mathcal{P}(A)$ is such that $U \cap G_A([p]) = U \cap G_A([\mathbf{1} - p])$, then $U \cap G_A([p]) = U \cap \{p\}' =: U(p)$.*

Proof. (i) We have

$$G_A([p]) = \{u \in G_A \mid [upu^{-1}] = [p]\} \text{ and } G_A([\mathbf{1} - p]) = \{u \in G_A \mid [u(\mathbf{1} - p)u^{-1}] = [\mathbf{1} - p]\},$$

so that clearly $G_A([p]) \cap G_A([\mathbf{1} - p]) \supseteq G_A \cap \{p\}'$. For the converse inclusion let $u \in G_A([p]) \cap G_A([\mathbf{1} - p])$ arbitrary. In particular $u \in G_A([p])$, whence $upu^{-1} \sim p$, which is equivalent to the fact that $(upu^{-1})p = p$ and $p(upu^{-1}) = upu^{-1}$. Consequently we have both

$$(2.1) \quad pu^{-1}p = u^{-1}p$$

and

$$(2.2) \quad pup = up.$$

On the other hand, since $u \in G_A([\mathbf{1} - p])$ as well, it follows that $(\mathbf{1} - p)u^{-1}(\mathbf{1} - p) = u^{-1}(\mathbf{1} - p)$ and $(\mathbf{1} - p)u(\mathbf{1} - p) = u(\mathbf{1} - p)$. The later equation is equivalent to $u - up - pu + pup = u - up$, that is, $pup = pu$. Then (2.2) implies that $up = pu$, that is, $u \in G(p)$.

(ii) This follows at once from part (i). \square

Remark 2.5. For instance, Proposition 2.4(ii) can be applied if the algebra A is equipped with an involution $a \mapsto a^*$ such that $p = p^*$, and $U = U_A := \{u \in G_A \mid u^{-1} = u^*\}$ is the corresponding unitary group. In this case, it follows by (2.1) and (2.2) that $up = pu$ whenever $u \in U_A \cap G_A([p])$, hence $U_A \cap G_A([p]) = U_A \cap G_A([\mathbf{1} - p]) = U_A \cap \{p\}' =: U_A(p)$.

For $q \in \mathcal{P}(A)$, put $\hat{q} := \mathbf{1} - q$ and $A^q := \{a \in A \mid \hat{q}aq = 0\}$. The following result is partly a counterpart, for algebras, of Proposition 2.4.

Proposition 2.6. *Assume that A is equipped with an involution and let $p \in \mathcal{P}(A)$ such that $p = p^*$. Then the following assertions hold:*

- (i) $uA^pu^{-1} = A^p$, for every $u \in U_A(p)$;
- (ii) $A^p \cap A^{\hat{p}} = \{p\}'$;
- (iii) $A^p + A^{\hat{p}} = A$;
- (iv) $(A^p)^* = A^{\hat{p}}$.

Proof. (i) This is readily seen.

(ii) Firstly, note that, for $a \in A$, we have $\hat{p}ap = pa\hat{p}$ if and only if $ap = pa$. Moreover, if $ap = pa$ then $\hat{p}ap = ap - pap = ap - ap = 0$ and analogously $pa\hat{p} = 0$. From this, the equality of the statement follows.

(iii) For every $a \in A$ and $q \in \mathcal{P}(A)$ we have $qa \in A^q$. Hence $a = pa + \hat{p}a \in A^p + A^{\hat{p}}$, as we wanted to show.

(iv) Take $a \in A^p$. Then $pap = ap$, that is, $pa^*p = pa^*$. Hence, $\hat{p}a^*\hat{p} = (a^* - pa^*)(1 - p) = a^* - pa^* - a^*p + a^*p = a^* - a^*p = a^*\hat{p}$. This means that $a^* \in A^{\hat{p}}$. Conversely, if $a \in A^{\hat{p}}$ then, as above, $pa^*p = a^*p$; that is, $a = (a^*)^*$ with $a^* \in A^p$. \square

3. HOMOGENEOUS COMPLEX STRUCTURES AND COMPLEXIFICATIONS

Definition 3.1. Let X be a Banach manifold. A *complexification* of X is a complex Banach manifold Y endowed with an anti-holomorphic involutive diffeomorphism $y \mapsto y^{-*}$ such that the fixed point submanifold $Y_0 = \{y \in Y \mid y = y^{-*}\}$ is diffeomorphic to X . \square

Assume from now on that A is a unital C^* -algebra. Then G_A is a Banach-Lie group whose Lie algebra coincides with A . The G_A -orbits in $\text{Gr}(A)$, obtained by the action β and equipped with the topology inherited from $\text{Gr}(A)$, are holomorphic Banach manifolds diffeomorphic to $G_A/G_A([p])$ (endowed with its quotient topology), see Theorem 2.2 in [DG02]. Also, the Grassmannian $\text{Gr}(A)$ can be described as the discrete union of these G_A -orbits, see [DG01] and Theorem 2.3 in [DG02]. Moreover, U_A is a Banach-Lie subgroup of G_A with the Lie algebra $\mathfrak{u}_A := \{a \in A \mid a^* = -a\}$. As it is well known, the complexification of \mathfrak{u}_A is A , via the decomposition $a = \{(a - a^*)/2\} + i\{(a + a^*)/2i\}$, ($a \in A$). Thus the conjugation of A is given by $a \mapsto \bar{a} := \{(a - a^*)/2\} - i\{(a + a^*)/2i\} = -a^*$. We seek for possible topological and/or differentiable relationships between the G_A -orbits and the U_A -orbits $U_A/U_A(p)$ in $\text{Gr}(A)$.

The above observations lead to the following result.

Theorem 3.2. Assume that A is a unital C^* -algebra, $p = p^* \in \mathcal{P}(A)$ and $\mathfrak{u}_A(p) := \mathfrak{u}_A \cap \{p\}'$. Let Ad_U denote the adjoint representation of U_A . Then the following assertions hold:

- (i) $\text{Ad}_U(u)A^p \subset A^p$, ($u \in U_A(p)$); $A^p \cap \overline{A^p} = \mathfrak{u}_A(p) + i\mathfrak{u}_A(p)$; $A^p + \overline{A^p} = A$.
- (ii) The manifold $U_A/U_A(p)$ has a U_A -invariant complex structure and is locally biholomorphic to $G_A/G_A([p])$.
- (iii) The manifold $G_A/G_A(p)$ endowed with the involutive diffeomorphism $aG_A(p) \mapsto (a^*)^{-1}G_A(p)$ is a complexification of $U_A/U_A(p)$.

Proof. It is clear that $\mathfrak{u}_A(p) + i\mathfrak{u}_A(p) = \{p\}'$. Now the first part of the statement follows by Proposition 2.6.

Also, there is a natural identification between $\mathfrak{u}_A/\mathfrak{u}_A(p)$ and the tangent space $T_{[p]}(U_A/U_A(p))$. Then assertion (ii) follows from Theorem 6.1 in [B06]. In fact, by assertion (i) it is readily seen that $\mathfrak{u}_A/\mathfrak{u}_A(p) \simeq A/A^p$ whence we obtain that $U_A/U_A(p)$ and $G_A/G_A([p])$ are locally diffeomorphic, and so $U_A/U_A(p)$ inherits the complex structure induced by $G_A/G_A([p])$.

For assertion (iii), it is easy to see that the mapping $aG_A(p) \mapsto (a^*)^{-1}G_A(p)$ is an anti-holomorphic diffeomorphism (which corresponds to the mapping $apa^{-1} \mapsto (a^*)^{-1}pa^*$, in terms of orbits). Then $aG_A(p) = (a^*)^{-1}G_A(p)$ if and only if $(a^*a)G_A(p) = G_A(p)$, that is, $(a^*a)p = p(a^*a)$. Using the functional calculus for C^* -algebras, we can pick $b := \sqrt{a^*a}$ in A and obtain $bp = pb$. Since $a^*a = b^2 = b^*b$ we have $(ab^{-1})^* = (b^{-1})^*a^* = (b^*)^{-1}a^* = ba^{-1} = (ab^{-1})^{-1}$ and therefore $u := ab^{-1} \in U_A$. Finally, $aG_A(p) = ubG_A(p) = uG_A(p) \equiv uU_A(p) \in U_A/U_A(p)$. \square

Remark 3.3. Since $G_A(p) \subset G_A([p])$, there exists the canonical projection $G_A/G_A(p) \rightarrow G_A/G_A([p])$. It is clear that its restriction to $U_A/U_A(p)$ becomes the identity map $U_A/U_A(p) \rightarrow U_A/U_A([p])$.

According to Proposition 2.4(i), idempotents like $apa^{-1} \equiv aG_A(p)$, for $a \in G_A$, can be alternatively represented as pairs $(a[p]a^{-1}, (a^*)^{-1}[p]a^*)$ so that the “orbit” $G_A/G_A(p)$ becomes a subset of the Cartesian product $G_A([p]) \times G_A([p])$. In this perspective, the preceding projection and diffeomorphism are given, respectively, by

$$(a[p]a^{-1}, (a^*)^{-1}[p]a^*) \mapsto a[p]a^{-1} \equiv (a[p]a^{-1}, a[p]a^{-1}), \quad G_A/G_A(p) \rightarrow G_A/G_A([p])$$

(so $(u[p]u^{-1}, u[p]u^{-1}) \mapsto upu^{-1} \equiv (u[p]u^{-1}, u[p]u^{-1})$, when $u \in U_A$) and

$$(a[p]a^{-1}, (a^*)^{-1}[p]a^*) \mapsto ((a^*)^{-1}[p]a^*, a[p]a^{-1}),$$

for every $a \in G_A$. \square

Remark 3.4. Theorem 3.2 relates to the setting of [BR07]. Namely, assume that B is a C^* -subalgebra of A , with $\mathbf{1} \in B \subseteq A$, for which there exist a conditional expectation $E: A \rightarrow B$ and a state $\varphi: A \rightarrow \mathbb{C}$ such that $\varphi \circ E = \varphi$. For $X \in \{A, B\}$, we denote by φ_X the state φ restricted to X . Let \mathcal{H}_X be the Hilbert space, and let $\pi_X: X \rightarrow \mathcal{B}(\mathcal{H}_X)$ be the corresponding cyclic representation obtained by the GNS construction applied to the state $\varphi_X: X \rightarrow \mathbb{C}$. Thus, \mathcal{H}_X is the completion of X/N_X with respect to the norm $\|y + N_X\|_\varphi := \varphi(y^*y)$, where $N_X := \{y \in X \mid \varphi(y^*y) = 0\}$. The representation π_X is then defined as the extension to \mathcal{H}_X of the left multiplication of X on X/N_X . Let P denote the orthogonal projection $P: \mathcal{H}_A \rightarrow \mathcal{H}_B$.

An equivalence relation can be defined in $G_A \times \mathcal{H}_B$ by setting that $(g_1, h_1) \sim (g_2, h_2)$ (with $g_1, g_2 \in G_A$, $h_1, h_2 \in \mathcal{H}_B$) if and only if there exists $w \in G_B$ such that $g_2 = g_1w$ and $h_2 = \pi_B(w^{-1})h_1$. The corresponding quotient space will be denoted by $G_A \times_{G_B} \mathcal{H}_B$, and the equivalence class in $G_A \times_{G_B} \mathcal{H}_B$ of an element $(g, h) \in G_A \times \mathcal{H}_B$ will be denoted by $[(g, h)]$. Define $U_A \times_{U_B} \mathcal{H}_B$ in an analogous fashion. Then the mappings $\Pi_G: [(g, h)] \mapsto gG_B$, $G_A \times_{G_B} \mathcal{H}_B \rightarrow G_A/G_B$ and $\Pi_U: [(u, h)] \mapsto uU_B$, $U_A \times_{U_B} \mathcal{H}_B \rightarrow U_A/U_B$ are vector bundles, Π_U being Hermitian, in fact. Moreover, Π_U admits a reproducing kernel K with the associated Hilbert space \mathcal{H}^K , formed by continuous sections of Π_U , such that the restriction of the GNS representation π_A to U_A can be realized on \mathcal{H}^K , see [BR07].

Let us apply the above theory to the case when, for a given unital C^* -algebra A , we take $B := \{p\}'$ in A , where $p = p^* \in \mathcal{P}(A)$. Then $E_p: a \mapsto pap + \hat{a}p\hat{a}$, $A \rightarrow B$ is a conditional expectation from A onto B . Let \mathcal{H} be a Hilbert space such that $A \hookrightarrow \mathcal{B}(\mathcal{H})$. Pick $x_0 \in p\mathcal{H}$ such that $\|x_0\| = 1$. Then $\varphi_0: A \rightarrow \mathbb{C}$, given by $\varphi_0(a) := (ax_0 \mid x_0)_\mathcal{H}$ for all $a \in A$, is a state of A such that $\varphi_0 \circ E_p = \varphi_0$. The GNS representation of A associated with φ_0 is as follows. Set $(a_1 \mid a_2)_0 := \varphi_0(a_2^*a_1) = (a_2^*a_1x_0 \mid x_0)_\mathcal{H} = (a_1x_0 \mid a_2x_0)_\mathcal{H}$ for every $a_1, a_2 \in A$. So $\varphi_0(a^*a) = \|a(x_0)\|^2$ for all $a \in A$, whence the null space of $(\cdot \mid \cdot)_0$ is $N_0 := \{a \in A : (a \mid a)_0 = 0\} = \{a \in A : a(x_0) = 0\}$. The norm $\|\cdot\|_0$ induced by $(\cdot \mid \cdot)_0$ on A/N_0 is given by $\|h\|_0 \equiv \|a + N_0\|_0 := \varphi_0(a^*a)^{1/2} = \|a(x_0)\|_\mathcal{H} = \|h\|_\mathcal{H}$ for every $h \in A(x_0) \subset \mathcal{H}$, where $a(x_0) = h \leftrightarrow a + N_0$. Hence \mathcal{H}_A is a closed subspace of \mathcal{H} such that $a\mathcal{H}_A \subset \mathcal{H}_A$ for every $a \in A$. Note that \mathcal{H}_A coincides with \mathcal{H} provided that we can choose x_0 in \mathcal{H} such that $A(x_0)$ is dense in \mathcal{H} . This will be of interest in Remark 4.8 below.

Analogously, we can consider the restriction of $(\cdot \mid \cdot)_0$ to B and proceed in the same way as above. Thus we obtain that the corresponding null space is $B \cap N_0$, that the norm in $B/(B \cap N_0)$ is that one of $p\mathcal{H}$ (so that one of \mathcal{H}), and that \mathcal{H}_B is a closed subspace of $p\mathcal{H}$ such that $b\mathcal{H}_B \subset \mathcal{H}_B$ for every $b \in B$. Also, $\mathcal{H}_B = p\mathcal{H}$ if x_0 can be chosen in $p\mathcal{H}$ and such that $B(x_0)$ is dense in $p\mathcal{H}$.

The representation $\pi_A: a \mapsto \pi(a)$, $A \rightarrow \mathcal{B}(\mathcal{H}_A)$ is the extension to \mathcal{H}_A of the left multiplication $\pi_A(a): a' + N_0 \mapsto (aa') + N_0$, $A/N_0 \rightarrow A/N_0$. Thus it satisfies $\pi_A(a' + N_0) = (aa') + N_0 \equiv a(a'x_0) = a(h)$, if $(a' + N_0) \leftrightarrow a(x_0) = h$. In other words, π_A is the inclusion operator (by restriction) from A into $\mathcal{B}(\mathcal{H}_A)$. Also, π_B is in turn the inclusion operator from B into $\mathcal{B}(\mathcal{H}_B)$.

Since $E_p(N_0) \subseteq N_0$, the expectation E_p induces a well-defined projection $P: A/N_0 \rightarrow B/(N_0 \cap B)$. On the other hand, $E_p(a^*a) - E_p(a)^*E_p(a) = pa^*\hat{p}ap + \hat{p}a^*pap \geq 0$ since $p, \hat{p} \geq 0$. Hence P extends once again as a bounded projection $P: \mathcal{H}_A \rightarrow \mathcal{H}_B$. Indeed, if $h = a(x_0)$ with $a \in A$, we have

$$P(h) \equiv P(a + N_0) = E(a) + (B \cap N_0) = E(a)(x_0) = (pa)(x_0) = p(h),$$

that is, $P = p|_{\mathcal{H}_A}$. \square

In the above setting, note that $U_B = U_A(p)$. Let $\Gamma(U_A/U_A(p), U_A \times_{U_A(p)} \mathcal{H}_B)$ be the section space of the bundle Π_U . The reproducing kernel associated with Π_U is given by $K_p(u_1U_A(p), u_2U_A(p))[(u_2, f_2)] := [(u_1, pu_1^{-1}u_2f_2)]$, for every $u_1, u_2 \in U_A$ and $f_2 \in \mathcal{H}_B$. The kernel K_p generates a Hilbert subspace \mathcal{H}^{K_p} of sections in $\Gamma(U_A/U_A(p), U_A \times_{U_A(p)} \mathcal{H}_B)$. Let $\gamma_p: \mathcal{H}_A \rightarrow \Gamma(U_A/U_A(p), U_A \times_{U_A(p)} \mathcal{H}_B)$ be the mapping defined by $\gamma_p(h)(uU_A(p)) := [(u, pu^{-1}h)]$ for every $h \in \mathcal{H}_A$ and $u \in U_A$. Then γ_p is injective and it intertwines the representation π_A of U_A on \mathcal{H}_A and the natural action of U_A on \mathcal{H}^{K_p} ; that is, the

diagram

$$(3.1) \quad \begin{array}{ccc} \mathcal{H}_A & \xrightarrow{u} & \mathcal{H}_A \\ \gamma_p \downarrow & & \downarrow \gamma_p \\ \mathcal{H}^{K_p} & \xrightarrow{\mu(u)} & \mathcal{H}^{K_p}, \end{array}$$

is commutative for all $u \in U_A$, where $\mu(u)F := uF(u^{-1} \cdot)$ for every $F \in \Gamma(U_A/U_A(p), U_A \times_{U_A(p)} \mathcal{H}_B)$. In fact $\gamma(uh)(vU_A(p)) := [(v, pv^{-1}uh)] = u[(u^{-1}v, pv^{-1}uh)] =: u\{\gamma(h)(u^{-1}vU_A(p))\}$ for all $u, v \in U_A$. See Theorem 5.4 of [BR07] for details in the general case. We next show that \mathcal{H}^{K_p} in fact consists of holomorphic sections.

Proposition 3.5. *Let A be a unital C^* -algebra, $p = p^* \in \mathcal{P}(A)$, and $B := \{p\}'$. Then the homogeneous Hermitian vector bundle $\Pi_U: U_A \times_{U_A(p)} \mathcal{H}_B \rightarrow U_A/U_A(p)$ is holomorphic, and the image of γ_p consists of holomorphic sections. Thus \mathcal{H}^{K_p} is a Hilbert space of holomorphic sections of Π_U .*

Proof. Let $u_0 \in U_A$. Then $\Omega_G := \{u_0g \mid g \in G_A, \|1 - g^{-1}\| < 1\}$ is open in G_A and contains u_0 , and similarly with $\Omega_U := \Omega_G \cap U_A$ in U_A .

It is readily seen that the mapping $\psi_0: [(u, f)] \mapsto (uU_A(p), E_p(u^{-1}u_0^{-1})^{-1}f)$, $\Pi_U^{-1}(\Omega_U) \rightarrow \Omega_U \times \mathcal{H}_B$ is a diffeomorphism, with inverse map $(uU_A(p), h) \mapsto [(uE_p(u^{-1}), h)]$ (this shows the local triviality of Π_U). Thus every point in the manifold $U_A \times_{U_A(p)} \mathcal{H}_B$ has an open neighborhood which is diffeomorphic to the manifold product $W \times \mathcal{H}_B$, where W is an open subset of $U_A/U_A(p)$. By Theorem 3.2, $U_A/U_A(p)$ is a complex homogeneous manifold and therefore the manifold $U_A \times_{U_A(p)} \mathcal{H}_B$ is locally complex, i.e., holomorphic. Also the bundle map Π_U is holomorphic.

On the other hand, for fixed $h \in \mathcal{H}_A$, the mapping $\sigma_0: gG_A([p]) \mapsto E_p(g^{-1}u_0^{-1})^{-1}pg^{-1}h$, $\Omega_G \rightarrow \mathcal{H}_B$ is holomorphic on Ω_G , so it defines a holomorphic function $\tilde{\sigma}_0: \Omega_G G_A([p]) \rightarrow \mathcal{H}_B$. By Theorem 3.2 the injection $j: U_A/U_A(p) \hookrightarrow G_A/G_A([p])$ is holomorphic, and so the restriction map $r := \tilde{\sigma}_0 \circ j$ is holomorphic around $u_0U_A(p)$. Since $\gamma(h) = \psi_0^{-1} \circ (I_{\Omega_U} \times r)$ around $u_0U_A(p)$, it follows that $\gamma(h)$ is (locally) holomorphic.

Finally, by applying Theorem 4.2 in [BR07] we obtain that K_p is holomorphic. \square

The starting point for the holomorphic picture given in Proposition 3.5 has been the fact that $U_A/U_A(p)$ enjoys a holomorphic structure induced by the one of $G_A/G([p])$, see Theorem 3.2. Such a picture can be made even more explicit if we have a global diffeomorphism $U_A/U_A(p) \simeq G_A/G_A([p])$. The prototypical example is to be found when A is the algebra of bounded operators on a complex Hilbert space. We examine this case more closely in the next section.

4. TAUTOLOGICAL UNIVERSAL VECTOR BUNDLES

Let us recall the specific definition and some properties of the Grassmannian manifold associated with a complex Hilbert space.

Notation 4.1. We shall use the standard notation $\mathcal{B}(\mathcal{H})$ for the C^* -algebra of bounded linear operators on the complex Hilbert space \mathcal{H} with the involution $T \mapsto T^*$. Let $\text{GL}(\mathcal{H})$ be the Banach-Lie group of all invertible elements of $\mathcal{B}(\mathcal{H})$, and $\text{U}(\mathcal{H})$ its Banach-Lie subgroup of all unitary operators on \mathcal{H} . Also,

- $\text{Gr}(\mathcal{H}) := \{\mathcal{S} \mid \mathcal{S} \text{ closed linear subspace of } \mathcal{H}\};$
- $\mathcal{T}(\mathcal{H}) := \{(\mathcal{S}, x) \in \text{Gr}(\mathcal{H}) \times \mathcal{H} \mid x \in \mathcal{S}\} \subseteq \text{Gr}(\mathcal{H}) \times \mathcal{H};$
- $\Pi_{\mathcal{H}}: (\mathcal{S}, x) \mapsto \mathcal{S}, \mathcal{T}(\mathcal{H}) \rightarrow \text{Gr}(\mathcal{H});$
- for every $\mathcal{S} \in \text{Gr}(\mathcal{H})$ we denote by $p_{\mathcal{S}}: \mathcal{H} \rightarrow \mathcal{S}$ the corresponding orthogonal projection.

\square

Remark 4.2. The objects introduced in Notation 4.1 have the following well known properties:

- (a) Both $\text{Gr}(\mathcal{H})$ and $\mathcal{T}(\mathcal{H})$ have structures of complex Banach manifolds, and $\text{Gr}(\mathcal{H})$ carries a natural (non-transitive) action of $\text{U}(\mathcal{H})$. (See Examples 3.11 and 6.20 in [U85], or Chapter 2 in [Du66].)

- (b) For every $\mathcal{S}_0 \in \text{Gr}(\mathcal{H})$ the corresponding connected component of $\text{Gr}(\mathcal{H})$ is the $\text{GL}(\mathcal{H})$ -orbit and is also the $\text{U}(\mathcal{H})$ -orbit of \mathcal{S}_0 , that is,

$$\begin{aligned} \text{Gr}_{\mathcal{S}_0}(\mathcal{H}) &= \{g\mathcal{S}_0 \mid g \in \text{GL}(\mathcal{H})\} = \{u\mathcal{S}_0 \mid u \in \text{U}(\mathcal{H})\} \\ &= \{\mathcal{S} \in \text{Gr}(\mathcal{H}) \mid \dim \mathcal{S} = \dim \mathcal{S}_0 \text{ and } \dim \mathcal{S}^\perp = \dim \mathcal{S}_0^\perp \simeq \text{U}(\mathcal{H})/(\text{U}(\mathcal{S}_0) \times \text{U}(\mathcal{S}_0^\perp))\}. \end{aligned}$$

(See Proposition 23.1 in [U85] or Lemma 4.3 below, alternatively.)

- (c) The mapping $\Pi_{\mathcal{H}}: \mathcal{T}(\mathcal{H}) \rightarrow \text{Gr}(\mathcal{H})$ is a holomorphic Hermitian vector bundle, and we call it the *universal (tautological) vector bundle* associated with the Hilbert space \mathcal{H} . Set $\mathcal{T}_{\mathcal{S}_0}(\mathcal{H}) := \{(\mathcal{S}, x) \in \mathcal{T}(\mathcal{H}) \mid \mathcal{S} \in \text{Gr}_{\mathcal{S}_0}(\mathcal{H})\}$. The vector bundle $\mathcal{T}_{\mathcal{S}_0}(\mathcal{H}) \rightarrow \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ obtained by restriction of $\Pi_{\mathcal{H}}$ to $\mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$ will be called here the *universal vector bundle at \mathcal{S}_0* . It is also Hermitian and holomorphic.

□

Property (b) in Remark 4.2 means that $U_A/U_A(p_{\mathcal{S}_0}) \simeq G_A/G_A([p_{\mathcal{S}_0}])$ for $A = \mathcal{B}(\mathcal{H})$. For the sake of clarification we now connect Notation 2.1 and Notation 4.1 in more detail. For $A = \mathcal{B}(\mathcal{H})$ we have $\text{Gr}(A) = \text{Gr}(\mathcal{H})$, and with this identification the action β of Lemma 2.2 corresponds to the natural action (so-called *collineation* action) of the group of invertible operators on \mathcal{H} upon the set of all closed linear subspaces of \mathcal{H} . The following lemma gives us the collineation orbits of $\text{Gr}(\mathcal{H})$ in terms of orbits of projections, and serves in particular to explain the property stated in Remark 4.2(b).

For short, denote $\mathcal{G} = \text{GL}(\mathcal{H})$ and $\mathcal{U} = \text{U}(\mathcal{H})$.

Lemma 4.3. *Let $\mathcal{S}_0 \in \text{Gr}(\mathcal{H})$. Then the following assertions hold.*

- (i) $\mathcal{G}([p_{\mathcal{S}_0}]) = \{g \in \mathcal{G} \mid g\mathcal{S}_0 = \mathcal{S}_0\}$ and $\mathcal{U}([p_{\mathcal{S}_0}]) = \mathcal{U}(p_{\mathcal{S}_0}) = \{u \in \mathcal{U} \mid u\mathcal{S}_0 = \mathcal{S}_0\}$.
- (ii) For every $g \in \mathcal{G}$ and $\mathcal{S} = g\mathcal{S}_0$ we have $\mathcal{S}^\perp = (g^*)^{-1}(\mathcal{S}_0^\perp)$.
- (iii) We have

$$\begin{aligned} \text{Gr}_{\mathcal{S}_0}(\mathcal{H}) &= \{g\mathcal{S}_0 \mid g \in \mathcal{G}\} \simeq \{[gp_{\mathcal{S}_0}g^{-1}] \mid g \in \mathcal{G}\} \\ &= \{u\mathcal{S}_0 \mid u \in \mathcal{U}\} \simeq \{up_{\mathcal{S}_0}u^{-1} \mid u \in \mathcal{U}\}. \end{aligned}$$

- (iv) We have

$$\mathcal{U}/\mathcal{U}(p_{\mathcal{S}_0}) \simeq \mathcal{G}/\mathcal{G}([p_{\mathcal{S}_0}]) \simeq \text{Gr}_{\mathcal{S}_0}(\mathcal{H}),$$

where the symbol “ \simeq ” means diffeomorphism between the respective differentiable structures, and that the differentiable structure of the quotient spaces is the one associated with the corresponding quotient topologies.

- (v) $\mathcal{G}/\mathcal{G}(p_{\mathcal{S}_0}) \simeq \{(a\mathcal{S}_0, (a^*)^{-1}\mathcal{S}_0) \mid a \in \mathcal{G}\}$ and the map $(a\mathcal{S}_0, (a^*)^{-1}\mathcal{S}_0) \mapsto ((a^*)^{-1}\mathcal{S}_0, a\mathcal{S}_0)$ is an involutive diffeomorphism on $\mathcal{G}/\mathcal{G}(p_{\mathcal{S}_0})$. Its set of fixed points is $\text{Gr}_{\mathcal{S}_0}(\mathcal{H}) \equiv \{(u\mathcal{S}_0, u\mathcal{S}_0) \mid u \in \mathcal{U}\}$.

Proof. (i) As shown in Proposition 2.4, an element g of \mathcal{G} belongs to $\mathcal{G}([p_{\mathcal{S}_0}])$ if and only if $p_{\mathcal{S}_0}g^{-1}p_{\mathcal{S}_0} = g^{-1}p_{\mathcal{S}_0}$ and $p_{\mathcal{S}_0}g p_{\mathcal{S}_0} = g p_{\mathcal{S}_0}$. From this, it follows easily that $g(\mathcal{S}_0) \subset \mathcal{S}_0$ and $g^{-1}(\mathcal{S}_0) \subset \mathcal{S}_0$, that is, $g(\mathcal{S}_0) = \mathcal{S}_0$. Conversely, if $g(\mathcal{S}_0) \subset \mathcal{S}_0$ then $(g p_{\mathcal{S}_0})(\mathcal{H}) \subset p_{\mathcal{S}_0}(\mathcal{H})$ whence $p_{\mathcal{S}_0}g p_{\mathcal{S}_0} = g p_{\mathcal{S}_0}$; similarly, $g^{-1}(\mathcal{S}_0) \subset \mathcal{S}_0$ implies that $p_{\mathcal{S}_0}g^{-1}p_{\mathcal{S}_0} = g^{-1}p_{\mathcal{S}_0}$. In conclusion, $\mathcal{G}([p_{\mathcal{S}_0}]) = \{g \in \mathcal{G} \mid g\mathcal{S}_0 = \mathcal{S}_0\}$.

Now, the above equality and Remark 2.5 imply that $\mathcal{U}([p_{\mathcal{S}_0}]) = \mathcal{U}(p_{\mathcal{S}_0}) = \{u \in \mathcal{U} \mid u\mathcal{S}_0 = \mathcal{S}_0\}$.

(ii) Let $x \in \mathcal{S}_0^\perp$, $y \in \mathcal{S}$. Then $((g^*)^{-1}(x) \mid y) = ((g^{-1})^*(x) \mid y) = (x \mid g^{-1}(y)) = 0$, so $(g^*)^{-1}(\mathcal{S}_0^\perp) \subset \mathcal{S}^\perp$. Take now $y \in \mathcal{S}^\perp$, $x = g^*(y)$ and $z \in \mathcal{S}_0$. Then $(x \mid z) = (g^*(y) \mid z) = (y \mid g(z)) = 0$, whence $x \in \mathcal{S}_0^\perp$ and therefore $y = (g^*)^{-1}(g^*y) = (g^*)^{-1}(x) \in (g^*)^{-1}(\mathcal{S}_0^\perp)$. In conclusion, $\mathcal{S}^\perp = (g^*)^{-1}(\mathcal{S}_0^\perp)$.

(iii) By (ii), we have $u(\mathcal{S}_0^\perp) = u(\mathcal{S}_0)^\perp$ for $u \in \mathcal{U}$. Thus $\mathcal{S} = u(\mathcal{S}_0)$ if and only if $\dim \mathcal{S} = \dim \mathcal{S}_0$ and $\dim \mathcal{S}^\perp = \dim \mathcal{S}_0^\perp$. Also, if $\mathcal{S} = u(\mathcal{S}_0)$ and $\mathcal{S}^\perp = u(\mathcal{S}_0^\perp)$, then $up_{\mathcal{S}_0} = p_{\mathcal{S}}u$, that is, $p_{\mathcal{S}} = up_{\mathcal{S}_0}u^{-1}$. Hence $\text{Gr}_{\mathcal{S}_0}(\mathcal{H}) = \{u\mathcal{S}_0 \mid u \in \mathcal{U}\} = \{\mathcal{S} \in \text{Gr}(\mathcal{H}) \mid \dim \mathcal{S} = \dim \mathcal{S}_0 \text{ and } \dim \mathcal{S}^\perp = \dim \mathcal{S}_0^\perp\} \simeq \{up_{\mathcal{S}_0}u^{-1} \mid u \in \mathcal{U}\}$.

Suppose now that $\mathcal{S} = g\mathcal{S}_0$ with $g \in \mathcal{G}$. Then $\dim \mathcal{S} = \dim \mathcal{S}_0$. By (ii) again, $\mathcal{S}^\perp = (g^*)^{-1}(\mathcal{S}_0^\perp)$ and so $\dim \mathcal{S}^\perp = \dim \mathcal{S}_0^\perp$. Hence $\mathcal{S} \in \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$. Finally, the bijective correspondence between $g\mathcal{S}_0$ and $g[p_{\mathcal{S}_0}]g^{-1}$ is straightforward.

- (iv) This is clearly a consequence of parts (iii) and (i) from above, and Theorem 2.2 in [DG02].

(v) For every $a \in \mathcal{G}$, the pairs $(a\mathcal{S}_0, (a^*)^{-1}\mathcal{S}_0)$ and $(a[p]a^{-1}, (a^*)^{-1}[p]a^*)$ are in a one-to-one correspondence, by part (iii) from above. Hence, this part (v) is a consequence of Remark 3.3. □

Parts (iv) and (v) of Lemma 4.3 tell us that the Grassmannian orbit $\text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ is a complex manifold which in turn admits a complexification, namely the orbit $\mathcal{G}/\mathcal{G}(p_{\mathcal{S}_0})$.

Remark 4.4. As said in Remark 4.2(b), every $\text{GL}(\mathcal{H})$ -orbit (and so every $\text{U}(\mathcal{H})$ -orbit) is a connected component of $\text{Gr}(\mathcal{H})$. Let us briefly discuss the connected components of $\text{Gr}(A)$ when A is an arbitrary unital C^* -algebra. Every element $g \in G_A$ has a unique polar decomposition $g = ua$ with $u \in U_A$ and $0 \leq a \in G_A$, hence there exists a continuous path $t \mapsto u \cdot ((1-t)\mathbf{1} + ta)$ in G_A that connects $u = u \cdot \mathbf{1}$ to $g = u \cdot a$. Thus every connected component of the G_A -orbit of $[p] \in \text{Gr}(A)$ contains at least one connected component of the U_A -orbit of $[p] \in \text{Gr}(A)$ for any idempotent $p \in \mathcal{P}(A)$. (Loosely speaking, the U_A -orbit of $[p]$ has more connected components than the G_A -orbit of $[p]$.) Example 7.13 in [PR87] shows that the C^* -algebra A of the continuous functions $S^3 \rightarrow M_2(\mathbb{C})$ has the property that there indeed exist G_A -orbits of elements $[p] \in \mathcal{P}(A)$ which are nonconnected.

If the unitary group U_A is connected (so that the invertible group G_A is connected), then all the U_A -orbits and the G_A -orbits in $\text{Gr}(A)$ are connected since continuous images of connected sets are always connected. On the other hand, as said formerly, the Grassmannian $\text{Gr}(A)$ is the discrete union of these G_A -orbits. Thus if the unitary group U_A is connected, then the connected components of $\text{Gr}(A)$ are precisely the G_A -orbits in $\text{Gr}(A)$. One important case of connected unitary group U_A is when A is a W^* -algebra (since every $u \in U_A$ can be written as $u = \exp(ia)$ for some $a = a^* \in A$ by the Borel functional calculus, hence the continuous path $t \mapsto \exp(ita)$ connects $\mathbf{1}$ to u in U_A). For W^* -algebras such that $\text{Gr}(A)$ is the discrete union of U_A -orbits, it is then clear that the G_A -orbits and the U_A -orbits coincide. This is the case if A is the algebra of bounded operators on a complex Hilbert space, as we have seen before. \square

The universal bundle $\mathcal{T}_{\mathcal{S}_0}(\mathcal{H}) \rightarrow \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ can be expressed as a vector bundle obtained from the so-called (principal) *Stiefel bundle* associated to $p_{\mathcal{S}_0} \leftrightarrow \mathcal{S}_0$, see [DG02]. A similar result holds, by replacing the Stiefel bundle with certain, suitable, of its sub-bundles. To see this, let us now introduce several mappings.

Put $p := p_{\mathcal{S}_0}$. We consider $\mathcal{G} \times_{\mathcal{G}([p])} \mathcal{S}_0$ and $\mathcal{U} \times_{\mathcal{U}(p)} \mathcal{S}_0$ as in Remark 3.4. Note that $g_1 \mathcal{S}_0 = g_2 \mathcal{S}_0$ and $g_1(h_1) = g_2(h_2)$ ($g_1, g_2 \in \mathcal{G}$, $h_1, h_2 \in \mathcal{S}_0$) if and only if $(g_1, h_1) \sim (g_2, h_2)$, via $w = g_1^{-1}g_2 \in \mathcal{G}([p])$, in $\mathcal{G} \times \mathcal{S}_0$. Hence, the mapping $v_{\mathcal{G}}: \mathcal{G} \times \mathcal{S}_0 \rightarrow \mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$ defined by $v_{\mathcal{G}}((g, h)) = (g\mathcal{S}_0, g(h))$ for every $(g, h) \in \mathcal{G} \times \mathcal{S}_0$, induces the usual (canonical) quotient map $\tilde{v}_{\mathcal{G}}: \mathcal{G} \times_{\mathcal{G}([p])} \mathcal{S}_0 \rightarrow \mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$. We denote by $\tilde{v}_{\mathcal{U}}$ the restriction map of $v_{\mathcal{G}}$ on $\mathcal{G} \times \mathcal{S}_0$. As above, the quotient mapping $\tilde{v}_{\mathcal{U}}: \mathcal{U} \times_{\mathcal{U}(p)} \mathcal{S}_0 \rightarrow \mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$ is well defined.

Since $\mathcal{U}(p) = \mathcal{U} \cap \mathcal{G}([p])$, the inclusion mapping $j: \mathcal{U} \times_{\mathcal{U}(p)} \mathcal{S}_0 \rightarrow \mathcal{G} \times_{\mathcal{G}([p])} \mathcal{S}_0$ is well defined. Note that $j = (\tilde{v}_{\mathcal{G}})^{-1} \circ \tilde{v}_{\mathcal{U}}$.

Finally, let $P_{\mathcal{G}}: \mathcal{G} \times_{\mathcal{G}([p])} \mathcal{S}_0 \rightarrow \mathcal{G}/\mathcal{G}([p])$ and $P_{\mathcal{U}}: \mathcal{U} \times_{\mathcal{U}(p)} \mathcal{S}_0 \rightarrow \mathcal{U}/\mathcal{U}(p)$ denote the vector bundles built in the standard way from the Stiefel sub-bundles $g \mapsto g\mathcal{G}([p]) \simeq g(\mathcal{S}_0)$, $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{G}([p]) \simeq \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ and $u \mapsto u\mathcal{U}(p) \simeq u(\mathcal{S}_0)$, $\mathcal{U} \rightarrow \mathcal{U}/\mathcal{U}(p) \simeq \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ respectively.

Proposition 4.5. *The following diagram is commutative in both sides, and the horizontal arrows are biholomorphic diffeomorphisms between the corresponding holomorphic structures*

$$\begin{array}{ccccc} \mathcal{T}_{\mathcal{S}_0}(\mathcal{H}) & \xrightarrow{(\tilde{v}_{\mathcal{U}})^{-1}} & \mathcal{U} \times_{\mathcal{U}(p)} \mathcal{S}_0 & \xrightarrow{j} & \mathcal{G} \times_{\mathcal{G}([p])} \mathcal{S}_0 \\ \Pi_{\mathcal{H}} \downarrow & & \downarrow P_{\mathcal{U}} & & \downarrow P_{\mathcal{G}} \\ \text{Gr}_{\mathcal{S}_0}(\mathcal{H}) & \xrightarrow{\simeq} & \mathcal{U}/\mathcal{U}(p) & \xrightarrow{\simeq} & \mathcal{G}/\mathcal{G}([p]) \end{array}$$

Proof. By construction, the mapping $\tilde{v}_{\mathcal{U}}$ is clearly one-to-one. Now we show that it is onto. Let $(\mathcal{S}, h) \in \mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$. This means that $h \in \mathcal{S}$ and that $\mathcal{S} = u\mathcal{S}_0$ for some $u \in \mathcal{U}$. Then $f := u^{-1}(h) \in \mathcal{S}_0$ and $h = u(f)$, whence $\tilde{v}_{\mathcal{U}}([(u, f)]) = (\mathcal{S}, h)$, where $[(u, f)]$ is the equivalence class of (u, f) in $\mathcal{U} \times_{\mathcal{U}(p)} \mathcal{S}_0$. Hence $\tilde{v}_{\mathcal{U}}$ is a bijective map.

Analogously, we have that $\tilde{v}_{\mathcal{G}}$ is bijective from $\mathcal{G} \times_{\mathcal{G}([p])} \mathcal{S}_0$ onto $\mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$ as well. As a consequence, $j = (\tilde{v}_{\mathcal{G}})^{-1} \circ \tilde{v}_{\mathcal{U}}$ is also bijective. It is straightforward to check that all the maps involved in the diagram above are smooth. \square

Example 4.6. By Proposition 4.5 one can show that the universal, tautological bundle $\Pi_{\mathcal{H}}: \mathcal{T}_{\mathcal{S}_0}(\mathcal{H}) \rightarrow \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ enters, as a canonical example, the framework outlined in Theorem 5.4 and Theorem 5.8 of [BR07]. To see this in terms of the bundle $\Pi_{\mathcal{H}}$ itself, first note that the commutant algebra $\{p_{\mathcal{S}_0}\}'$ of $p_{\mathcal{S}_0}$ coincides with the Banach subalgebra B of A formed by the operators T such that $T(\mathcal{S}_0) \subset \mathcal{S}_0$, $T(\mathcal{S}_0^\perp) \subset \mathcal{S}_0^\perp$. (It is straightforward to check directly on B that it is stable under the adjoint operation, so that B is a C^* -subalgebra of A , as it had to be.) Put $p = p_{\mathcal{S}_0}$. From Lemma 4.3, $u \in \mathcal{U}([p])$ if and only if $u\mathcal{S}_0 = \mathcal{S}_0$. Hence $u \in \mathcal{U}(p) = \mathcal{U}([p]) \cap \mathcal{U}([1-p])$ if and only if $u\mathcal{S}_0 = \mathcal{S}_0$ and $u\mathcal{S}_0^\perp = \mathcal{S}_0^\perp$, that is, $\mathcal{U}(p) = \mathcal{U}_A \cap B = \mathcal{U}_B$.

Similarly to what has been done in Remark 3.4, let $E_p: A \rightarrow B$ denote the canonical expectation associated to the tautological bundle at \mathcal{S}_0 ; that is, $E_p(T) := pTp + \hat{p}T\hat{p}$ for every $T \in A$. Also, for a fixed $x_0 \in \mathcal{S}_0$ such that $\|x_0\| = 1$, let $\varphi: A \rightarrow \mathbb{C}$ be the state of A given by $\varphi_0(T) := (Tx_0 | x_0)_{\mathcal{H}}$. Then $\varphi_0 \circ E_p = \varphi_0$. Since the mappings $T \mapsto T(x_0)$, $\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{H}$ and $T \mapsto T(x_0)$, $B \rightarrow \mathcal{S}_0$ are surjective, we obtain that $\mathcal{H}_A = \mathcal{H}$ and $\mathcal{H}_B = \mathcal{S}_0$ in the GNS construction associated with $A = \mathcal{B}(\mathcal{H})$, B and φ_0 . Moreover, in this case, π_A coincides with the identity operator and the extension $P: \mathcal{H}_A \rightarrow \mathcal{H}_B$ of E_p is $P = p$. Denote by $p_1, p_2: \text{Gr}(\mathcal{H}) \times \text{Gr}(\mathcal{H}) \rightarrow \text{Gr}(\mathcal{H})$ the natural projections and define

$$Q_{\mathcal{H}}: \text{Gr}(\mathcal{H}) \times \text{Gr}(\mathcal{H}) \rightarrow \text{Hom}(p_2^*(\Pi_{\mathcal{H}}), p_1^*(\Pi_{\mathcal{H}}))$$

by

$$Q_{\mathcal{H}}(\mathcal{S}_1, \mathcal{S}_2) = (p_{\mathcal{S}_1})|_{\mathcal{S}_2}: \mathcal{S}_2 \rightarrow \mathcal{S}_1$$

whenever $\mathcal{S}_1, \mathcal{S}_2 \in \text{Gr}(\mathcal{H})$. This mapping $Q_{\mathcal{H}}$ is called the *universal reproducing kernel* associated with the Hilbert space \mathcal{H} . In fact, for $\mathcal{S}_1, \dots, \mathcal{S}_n \in \text{Gr}(\mathcal{H})$ and $x_j \in \mathcal{S}_j$ ($j = 1, \dots, n$),

$$\sum_{j,l=1}^n (Q_{\mathcal{H}}(\mathcal{S}_l, \mathcal{S}_j)x_j | x_l)_{\mathcal{H}} = \sum_{j,l=1}^n (p_{\mathcal{S}_l}x_j | x_l)_{\mathcal{H}} = \sum_{j,l=1}^n (x_j | x_l)_{\mathcal{H}} = \left(\sum_{j=1}^n x_j | \sum_{l=1}^n x_l \right)_{\mathcal{H}} \geq 0,$$

so $Q_{\mathcal{H}}$ is certainly a reproducing kernel in the sense of [BR07]. \square

Using Proposition 4.5 and Example 4.6 we get the following special case of Theorem 5.8 in [BR07].

Corollary 4.7. *For a complex Hilbert space \mathcal{H} , the action of \mathcal{U} on \mathcal{H} can be realized as the natural action of \mathcal{U} on a Hilbert space of holomorphic sections from $\text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ into \mathcal{H} , such a realization being implemented by $\gamma(uh) = u \gamma(h)u^{-1}$, for every $h \in \mathcal{H}$, $u \in \mathcal{U}$.*

Proof. If $\mathcal{S} \in \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$, there exists $u \in \mathcal{U}$ such that $u\mathcal{S}_0 = \mathcal{S}$ and then $p_{\mathcal{S}} = up_{\mathcal{S}_0}u^{-1}$. Thus for all $u_1, u_2 \in \mathcal{U}$ and $x_1, x_2 \in \mathcal{S}_0$ we have $Q_{\mathcal{H}}(u_1\mathcal{S}_0, u_2\mathcal{S}_0)(u_2x_2) = p_{u_1\mathcal{S}_0}(u_2x_2) = u_1p_{\mathcal{S}_0}(u_1^{-1}u_2x_2)$. This formula shows that for every connected component $\text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ the restriction of $Q_{\mathcal{H}}$ to $\text{Gr}_{\mathcal{S}_0}(\mathcal{H}) \times \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ is indeed a special case of the reproducing kernels considered in Remark 3.4. For every $h \in \mathcal{H}$, the mapping $\gamma_{p_{\mathcal{S}_0}}(h): \text{Gr}_{\mathcal{S}_0}(\mathcal{H}) \rightarrow \mathcal{T}_{\mathcal{S}_0}(\mathcal{H})$ which corresponds to $Q_{\mathcal{H}}$ can be identified to the holomorphic map $u\mathcal{S}_0 \mapsto upu^{-1}h$, $\text{Gr}_{\mathcal{S}_0}(\mathcal{H}) \rightarrow \mathcal{H}$. Then the conclusion follows by using the diffeomorphism $\mathcal{U}/\mathcal{U}(p) \simeq \mathcal{G}/\mathcal{G}([p]) \simeq \text{Gr}_{\mathcal{S}_0}(\mathcal{H})$ of Lemma 4.3, together with Proposition 4.5. \square

Remark 4.8. Assume again the situation where A and B are arbitrary C^* -algebras, B is a C^* -subalgebra of A , with unit $\mathbf{1} \in B \subseteq A$, $E: A \rightarrow B$ is a conditional expectation, and $\varphi: A \rightarrow \mathbb{C}$ is a state such that $\varphi \circ E = \varphi$. With the same notations as in Remark 3.4, take $x_0 := \mathbf{1} + N_B \in B/N_B \subset A/N_A$. It is well known that x_0 is a cyclic vector of π_X , for $X \in \{A; B\}$: let $h \in \mathcal{H}_X$ such that $0 = (\pi(c)x_0 | h)_{\mathcal{H}_X} \equiv (c + N_X | h)_{\mathcal{H}_X}$ for all $c \in X$; since X/N_X is dense in \mathcal{H}_X we get $0 = (h | h)_{\mathcal{H}_X} = \|h\|^2$, that is, $h = 0$. Thus $\pi_X(X)x_0$ is dense in \mathcal{H}_X .

Inspired by [AS94], we now consider the C^* -subalgebra \mathfrak{A} of $\mathcal{B}(\mathcal{H}_A)$ generated by $\pi_A(A)$ and p , where p is the orthogonal projection from \mathcal{H}_A onto \mathcal{H}_B . Set $\mathfrak{B} := \mathfrak{A} \cap \{p\}'$. Clearly, the GNS procedure is applicable to $\mathfrak{B} \subset \mathfrak{A} \subset \mathcal{B}(\mathcal{H}_A)$, for the expectation $E_p: \mathfrak{A} \rightarrow \mathfrak{B}$ and state φ_0 defined by x_0 , as we have done in Remark 3.4. Then $\pi_A(A)(x_0) \subset \mathfrak{A}(x_0) \subset \mathcal{H}_A$ and $\pi_A(B)(x_0) \subset \mathfrak{B}(x_0) \subset \mathcal{H}_B$, whence, by the choice of x_0 , we obtain that $\overline{\mathfrak{A}(x_0)} = \mathcal{H}_A$ and $\overline{\mathfrak{B}(x_0)} = \mathcal{H}_B$. Thus we have that $\mathcal{H}_{\mathfrak{A}} = \mathcal{H}_A$ and $\mathcal{H}_{\mathfrak{B}} = \mathcal{H}_B$.

According to former discussions there are two (composed) commutative diagrams, namely

$$(4.1) \quad \begin{array}{ccccccc} G_A \times_{G_B} \mathcal{H}_B & \xrightarrow{\pi_A \tilde{\times} I} & G_{\mathfrak{A}} \times_{G_{\mathfrak{A}}(p)} \mathcal{H}_B & \longrightarrow & G_{\mathfrak{A}} \times_{G_{\mathfrak{A}}([p])} \mathcal{H}_B & \xrightarrow{j \tilde{\times} I} & \mathcal{G} \times_{\mathcal{G}([p])} \mathcal{H}_B \\ \Pi_G \downarrow & & \downarrow \Pi_{G_{\mathfrak{A}}} & & \downarrow & & \downarrow \Pi_{\mathcal{H}_B} \\ G_A/G_B & \xrightarrow{\tilde{\pi}_A} & G_{\mathfrak{A}}/G_{\mathfrak{A}}(p) & \longrightarrow & G_{\mathfrak{A}}/G_{\mathfrak{A}}([p]) & \xrightarrow{\tilde{j}} & \mathcal{G}/\mathcal{G}([p]) \end{array}$$

and

$$(4.2) \quad \begin{array}{ccccccc} U_A \times_{U_B} \mathcal{H}_B & \xrightarrow{\pi_A \tilde{\times} I} & U_{\mathfrak{A}} \times_{U_{\mathfrak{A}}(p)} \mathcal{H}_B & \xrightarrow{j \tilde{\times} I} & \mathcal{U} \times_{\mathcal{U}(p)} \mathcal{H}_B & \xrightarrow{\simeq} & \mathcal{T}_{\mathcal{H}_B}(\mathcal{H}_A) \\ \Pi_U \downarrow & & \downarrow \Pi_{U_{\mathfrak{A}}} & & \downarrow \Pi_{\mathcal{U}} & & \downarrow \Pi_{\mathcal{H}_B} \\ U_A/U_B & \xrightarrow{\tilde{\pi}_A} & U_{\mathfrak{A}}/U_{\mathfrak{A}}(p) & \xrightarrow{\tilde{j}} & \mathcal{U}/\mathcal{U}(p) & \xrightarrow{\simeq} & \text{Gr}_{\mathcal{H}_B}(\mathcal{H}_A) \end{array}$$

(where the meaning of the arrows is clear). We suggest to call $\Pi_G: G_A \times_{G_B} \mathcal{H}_B \rightarrow G_A/G_B$ and $\Pi_U: U_A \times_{U_B} \mathcal{H}_B \rightarrow U_A/U_B$ the *GNS vector bundle* and the *unitary GNS vector bundle*, respectively, for data $E: A \rightarrow B$ and $\varphi: A \rightarrow \mathbb{C}$. Following the terminology used in [AS94], [ALRS97] for the maps $G_A/G_B \rightarrow G_{\mathfrak{A}}/G_{\mathfrak{A}}(p)$, $U_A/U_B \rightarrow U_{\mathfrak{A}}/U_{\mathfrak{A}}(p)$, we could refer to the left sub-diagrams of (4.1) and (4.2) as the *basic* vector bundle representations of Π_G and Π_U , respectively. Since $\mathcal{H}_{\mathfrak{A}} = \mathcal{H}_A$ and $\mathcal{H}_{\mathfrak{B}} = \mathcal{H}_B$, the process to construct such “basic” objects, of Grassmannian type, is stationary. Also, since there is another way to associate Grassmannians to the GNS and unitary GNS bundles, which is that one of considering the tautological bundle of \mathcal{H}_A (see the right diagrams in (4.1), (4.2)), we might call $G_{\mathfrak{A}} \times_{G_{\mathfrak{A}}(p)} \mathcal{H}_B \rightarrow G_{\mathfrak{A}}/G_{\mathfrak{A}}([p])$ the *minimal* Grassmannian vector bundle, and call $\mathcal{T}_{\mathcal{H}_B}(\mathcal{H}_A) \rightarrow \text{Gr}_{\mathcal{H}_B}(\mathcal{H}_A)$ the *universal* Grassmannian vector bundle, associated with data $E: A \rightarrow B$ and $\varphi: A \rightarrow \mathbb{C}$. In the unitary case, we should add the adjective “unitary” to both bundles.

Note that the vector bundles $\mathcal{G} \times_{\mathcal{G}([p])} \mathcal{H}_B \rightarrow \mathcal{G}/\mathcal{G}([p])$ and $\mathcal{T}_{\mathcal{H}_B}(\mathcal{H}_A) \rightarrow \text{Gr}_{\mathcal{H}_B}(\mathcal{H}_A)$ are isomorphic. In this sense, both diagrams (4.1) and (4.2) “converge” towards the tautological bundle for \mathcal{H}_A . Let us remark that (4.1) is holomorphic, and everything in (4.2) is holomorphic with the only possible exception of the bundle Π_U . On the other hand, we have that $G_{\mathfrak{A}}/G_{\mathfrak{A}}(p)$ and $\mathcal{G}/\mathcal{G}(p)$ are complexifications of $U_{\mathfrak{A}}/U_{\mathfrak{A}}(p)$ and $\mathcal{U}/\mathcal{U}(p)$ respectively, on account of Remark 3.3 and Lemma 4.3. Note in passing that the fact that G_A/G_B is such a complexification implies interesting properties of metric nature in the differential geometry of U_A/U_B , see [ALRS97].

The above considerations strongly suggest to investigate the relationships between (4.1) and (4.2) in terms of holomorphy and geometric realizations. In this respect, note that the commutativity of (4.2) corresponds, on the level of reproducing kernels, with the equality

$$(\pi_A \tilde{\times} I) \circ K(u_1 U_B, u_2 U_B) Q_{\mathcal{H}_B}(\pi_A(u_1) \mathcal{U}(p), \pi_A(u_2) \mathcal{U}(p)) \circ (\pi_A \tilde{\times} I)$$

for all $u_1, u_2 \in U_A$ (where the holomorphy supplied by $Q_{\mathcal{H}_B}$ appears explicitly). From this, a first candidate to reproducing kernel on G_A/G_B , in order to obtain a geometric realization of π_A on G_A , would be defined by

$$K(g_1 G_B, g_2 G_B)[(g_2, f)] := [(g_1, p(\pi_A(g_1^{-1}) \pi_A(g_2) f))]$$

for every $g_1, g_2 \in G_A$ and $f \in \mathcal{H}_B$. Nevertheless, since the elements g_1, g_2 are not necessarily unitary, it is readily seen that the kernel K so defined need not be definite-positive in general. There is also the problem of the existence of a suitable structure of Hermitian type in Π_G .

It would be interesting to have a theory of bundles $G_A \times_{G_B} \mathcal{H}_B \rightarrow G_A/G_B$ and kernels K taking into account natural involutive diffeomorphisms in G_A/G_B , which would allow to incorporate those bundles to a framework containing as a special case the one established in [BR07]. This will be the subject of a forthcoming paper by the authors. \square

5. ALMOST HYPERCOMPLEX STRUCTURES ASSOCIATED WITH GRASSMANN MANIFOLDS

The following definition provides the infinite-dimensional version of the terminology of quaternionic structures on finite-dimensional manifolds; see for instance subsection 2.5 in [AM96].

Definition 5.1. Let Y be a Banach manifold. An *almost hypercomplex structure* on Y is a pair of almost complex structures $J_1, J_2: TY \rightarrow TY$ satisfying $J_1 J_2 = -J_2 J_1$. \square

Remark 5.2. Let $\mathbb{H} = \mathbb{R} + \mathbb{R}i_1 + \mathbb{R}i_2 + \mathbb{R}i_3$ be the quaternion field with the imaginary units $i_1, i_2, i_3 \in \mathbb{H}$ satisfying $(i_1)^2 = (i_2)^2 = (i_3)^2 = -1$, $i_1 i_2 = -i_2 i_1 = i_3$, $i_2 i_3 = -i_3 i_2 = i_1$, and $i_3 i_1 = -i_1 i_3 = i_2$. In the setting of Definition 5.1, it is easy to see that the hypercomplex structure of Y gives rise to a fiberwise linear action of \mathbb{H} on TY by $i_1 \cdot v = J_1 v$, $i_2 \cdot v = J_2 v$ and $i_3 \cdot v = J_1 J_2 v$ for every $v \in TY$. Thus for every $y \in Y$ the tangent space $T_y Y$ has a natural structure of quaternionic vector space. \square

In the following statement we need the notion of spray on Banach manifolds in the sense of [L01].

Theorem 5.3. Assume that X is an almost complex Banach manifold. Then the following assertions hold:

- (i) There exists a natural correspondence from the sprays on X to the almost hypercomplex structures on TX .
- (ii) If there exist a unital C^* -algebra A and a projection $p = p^2 = p^* \in A$ such that $X = U_A/U_A(p)$, then the almost hypercomplex structure associated with the natural spray on X induces an almost hypercomplex structure on the complexification $G_A/G_A(p)$.

Proof. (i) Denote $Y = TX$ and $\pi: TX \rightarrow X$ the natural projection, and consider the commutative diagram

$$\begin{array}{ccc} \pi^*(TX) & \longrightarrow & TX \\ \pi^*(\pi) \downarrow & & \downarrow \pi \\ Y & \xrightarrow{\pi} & X \end{array}$$

where the left-hand vertical arrow is the pull-back of the right-hand vertical arrow by $\pi: Y \rightarrow X$. Assume that we have got the covariant derivative associated with some spray on X . It then follows by the tensorial splitting theorem (Theorem 4.3 in Chapter X of [L01]) that there exists an isomorphism

$$(5.1) \quad TY \simeq \pi^*(TX) \oplus_{TX} \pi^*(TX)$$

of vector bundles over TX . Note that the fiber of $\pi^*(TX)$ over any $y \in Y$ is

$$(\pi^*(TX))_y = \{(y, z) \in Y \times TX \mid \pi(y) = \pi(z)\} \simeq \pi^{-1}(\pi(y))$$

hence the fiber of the Whitney sum $\pi^*(TX) \oplus_{TX} \pi^*(TX)$ over $y \in Y = TX$ is

$$(\pi^*(TX) \oplus_{TX} \pi^*(TX))_y \{ (y_1, y_2) \in TX \times TX \mid \pi(y_1) = \pi(y_2) = \pi(y) \} \simeq T_{\pi(y)} X \times T_{\pi(y)} X.$$

By taking into account the isomorphism (5.1) we can now define two almost complex structures on Y by

$$(5.2) \quad (y_1, y_2) \mapsto (-y_2, y_1), \quad TY \xrightarrow{J_1} TY,$$

and

$$(5.3) \quad (y_1, y_2) \mapsto (iy_1, -iy_2), \quad TY \xrightarrow{J_2} TY.$$

Note that for every pair $(y_1, y_2) \in \pi^*(TX) \oplus_{TX} \pi^*(TX) \simeq TY$ we have $J_1 J_2 (y_1, y_2) = J_1 (iy_1, -iy_2) = (iy_2, iy_1)$ and $J_2 J_1 (y_1, y_2) = J_2 (-y_2, y_1) = (-iy_2, -iy_1)$. Hence $J_1 J_2 = -J_2 J_1$, and thus the pair of almost complex structures J_1, J_2 defines an almost hypercomplex structure on $Y = TX$.

(ii) Now assume that $X = U_A/U_A(p)$ as in the statement. This is a complex homogeneous space by Theorem 3.2. The natural connection on this Grassmann manifold is the connection associated with the conditional expectation

$$E: A \rightarrow B, \quad E(a) = pap + (\mathbf{1} - p)a(\mathbf{1} - p),$$

where $B = \{a \in A \mid ap = pa\}$. Recall that this conditional expectation induces a connection in the principal bundle $U_A \rightarrow U_A/U_A(p)$ (see [ALRS97] and [G06]). On the other hand, if we denote

$\mathfrak{p} = \{a \in \text{Ker } E \mid a^* = -a\}$, then $U_A(p)$ acts upon \mathfrak{p} by means of the adjoint action and it is well known that there exists an isomorphism of vector bundles $U_A \times_{U_A(p)} \mathfrak{p} \simeq TX$ over $U_A/U_A(p) = X$. In particular the tangent bundle of X is a vector bundle associated with the principal bundle $U_A \rightarrow U_A/U_A(p)$. Thus we get a linear connection on the vector bundle $TX \rightarrow X$ which is associated with a connection map (or connector) $TTX \rightarrow TX$, and the latter map gives rise to a spray on X by means of the Christoffel symbols. (See [L01] and subsections 37.24–27 in [KM97] for more details.) Now assertion (i) shows that there exists an almost hypercomplex structure on TX associated with the spray we got.

To complete the proof we have to show that there exists a projection $\pi: G_A/G_A(p) \rightarrow X$ making $G_A/G_A(p)$ into a vector bundle which is isomorphic to the tangent bundle $TX \rightarrow X$. Recall from the above reasonings that $TX \simeq U_A \times_{U_A(p)} \mathfrak{p}$ as vector bundles over $U_A/U_A(p)$. Now define the mapping $(u, a) \mapsto u \exp(ia)G_A(p)$, $U_A \times \mathfrak{p} \rightarrow G_A/G_A(p)$. It is straightforward to check that this induces an injective mapping $U_A \times_{U_A(p)} \mathfrak{p} \rightarrow G_A/G_A(p)$, which is actually a diffeomorphism as a consequence of Theorem 8 in [PR94]. This makes $G_A/G_A(p)$ into a vector bundle isomorphic to TX over X , and the proof ends. \square

Remark 5.4. The correspondence between affine connections on finite dimensional manifolds and almost (hyper)complex structures goes back to [H60] and [Do62]. See [BG96], [BG98], and [Bi03] for more recent advances. \square

Remark 5.5. Let us note another general way to construct almost hypercomplex structures associated with the infinite-dimensional complex Grassmann manifolds. Quite generally, assume that X is an almost complex Banach manifold. If we denote by \overline{X} the complex-conjugate manifold of X , then the direct product $X \times \overline{X}$ is a complexification of X and has a natural almost hypercomplex structure.

This fact was noted in the paper [D81] in the case of finite-dimensional manifolds and can be proved in the general case as follows. Let $I: TX \rightarrow TX$ be the almost complex structure of M . Then \overline{X} is just the underlying real analytic manifold of X thought of as an almost complex manifold with respect to the almost complex structure $-I: TX \rightarrow TX$. Let us denote by $\theta: X \rightarrow \overline{X}$ the identity mapping, which is an anti-holomorphic mapping. Also denote $Z = X \times \overline{X}$. Now consider the direct product almost complex structure of Z ,

$$J_1 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} : TZ \rightarrow TZ$$

and define

$$J_2 = \begin{pmatrix} 0 & T\theta \\ -(T\theta)^{-1} & 0 \end{pmatrix} : TZ \rightarrow TZ.$$

It is straightforward to check that $(J_1)^2 = (J_2)^2 = -\text{id}_{TZ}$ and $J_1 J_2 = -J_2 J_1$, where the latter equality follows by the fact that θ is antiholomorphic. \square

Acknowledgment. We wish to thank Professor Karl-Hermann Neeb and Professor Radu Pantilie for pointing out to us some pertinent references. The present research was begun during the first-named author's visit at the Department of Mathematics of the University of Zaragoza with financial support from Proyecto MTM2004-03036, DGI-FEDER, of the MCYT, Spain. Our research was further developed during the visit of the second-named author's visit at the Institute of Mathematics "Simion Stoilow" of the Romanian Academy with support from the aforementioned grant and from the Softwin Group. Partial financial support from Proyecto E-64 of the DG Aragón, Spain, and from Grant 2-CEX06-11-22/25.07.06 of the Romanian Government is also acknowledged.

REFERENCES

- [AM96] D.V. Alekseevsky, S. Marchiafava, Quaternionic structures on a manifold and subordinated structures, *Ann. Mat. Pura Appl. (4)* **171** (1996), 205–273.
- [ALRS97] E. Andruchow, A. Larotonda, L. Recht, D. Stojanoff, Infinite dimensional homogeneous reductive spaces and finite index conditional expectations, *Illinois J. Math.* **41** (1997), 54–76.
- [AS94] E. Andruchow, D. Stojanoff, Geometry of conditional expectations and finite index, *Int. J. Math.* **5** (1994), no. 2, 169–178.

- [B06] D. Beltiță, *Smooth Homogeneous Structures in Operator Theory*, Monogr. and Surveys in Pure and Applied Math., 137. Chapman & Hall/CRC Press, Boca Raton-London-New York-Singapore, 2006.
- [BR07] D. Beltiță, T.S. Ratiu, Geometric representation theory for unitary groups of operator algebras, *Adv. Math.* **208** (2007), no. 1, 299–317.
- [BN05] W. Bertram, K.-H. Neeb, Projective completions of Jordan pairs, Part II. Manifold structures and symmetric spaces, *Geom. Dedicata* **112** (2005), 73–113.
- [Bi03] R. Bielawski, Complexification and hypercomplexification of manifolds with a linear connection, *Internat. J. Math.* **14** (2003), no. 8, 813–824.
- [Bi04] R. Bielawski, Prescribing Ricci curvature on complexified symmetric spaces, *Math. Res. Lett.* **11** (2004), no. 4, 435–441.
- [BG96] O. Biquard, P. Gauduchon, La métrique hyperkählérienne des orbites coadjointes de type symétrique d’un groupe de Lie complexe semi-simple, *C. R. Acad. Sci. Paris Sér. I Math.* **323** (1996), no. 12, 1259–1264.
- [BG98] O. Biquard, P. Gauduchon, Géométrie hyperkählérienne des espaces hermitiens symétriques complexifiés, *Sémin. Théor. Spectr. Géom.*, vol. 16 (1997–1998), Univ. Grenoble I, Saint-Martin-d’Hères, 1998, pp. 127–173.
- [Bo57] R. Bott, Homogeneous vector bundles, *Ann. Math.* **66** (1957), no. 2, 203–248.
- [CPR90] G. Corach, H. Porta, L. Recht, Differential geometry of systems of projections in Banach algebras, *Pacific J. Math.* **143**(1990), no. 2, 209–228.
- [D81] G. Dedene, Complexifications of Lorentz- and symplectic structures, *Simon Stevin* **55** (1981), no. 3, 149–167.
- [Do62] P. Dombrowski, On the geometry of the tangent bundle, *J. Reine Angew. Math.* **210**(1962), 73–88.
- [Du66] A. Douady, Le problème des modules pour les sous-espaces analytiques compacts d’un espace analytique donné, *Ann. Inst. Fourier (Grenoble)* **16** (1966), no. 1, 1–95.
- [DG01] M.J. Dupré, J.F. Glazebrook, The Stiefel bundle of a Banach algebra, *Integral Equations Operator Theory* **41** (2001), no. 3, 264–287.
- [DG02] M.J. Dupré, J.F. Glazebrook, Holomorphic framings for projections in a Banach algebra, *Georgian Math. J.* **9** (2002), no. 3, 481–494.
- [G06] J.E. Galé, Geometría de órbitas de representaciones de grupos y álgebras promediabiles, *Rev. R. Acad. Cienc. Exactas Fís. Quím. Nat. Zaragoza (2)* **61** (2006), 7–46.
- [H60] C. Hsu, On some structures which are similar to the quaternion structure, *Tôhoku Math. J. (2)* **12** (1960), 403–428.
- [KM97] A. Kriegl, P.W. Michor, *The Convenient Setting of Global Analysis*, Mathematical Surveys and Monographs 53, American Mathematical Society, Providence, RI, 1997.
- [L01] S. Lang, *Fundamentals of Differential Geometry* (corrected second printing), Graduate Texts in Mathematics, 191. Springer-Verlag, New-York, 2001.
- [LS91] L. Lempert, R. Szöke, Global solutions of the homogeneous complex Monge-Ampère equation and complex structures on the tangent bundle of Riemannian manifolds, *Math. Ann.* **290** (1991), no. 4, 689–712.
- [MS98] M. Martin, N. Salinas, The canonical complex structure of flag manifolds in a C^* -algebra. In: H. Bercovici (ed.) et al., *Nonselfadjoint Operator Algebras, Operator Theory, and Related Topics. The Carl M. Pearcy Anniversary Volume on the Occasion of His 60th Birthday*, Birkhäuser Verlag, Oper. Theory Adv. Appl., 104, Basel, 1998, pp. 173–187.
- [MR92] L.E. Mata-Lorenzo, L. Recht, Infinite-dimensional homogeneous reductive spaces, *Acta Cient. Venezolana* **43**(1992), no. 2, 76–90.
- [PR87] H. Porta, L. Recht, Spaces of projections in a Banach algebra, *Acta Cient. Venezolana* **38** (1987), no. 4, 408–426.
- [PR94] H. Porta, L. Recht, Conditional expectations and operator decompositions, *Ann. Global Anal. Geom.* **12** (1994), no. 4, 335–339.
- [S88] N. Salinas, The Grassmann manifold of a C^* -algebra, and Hermitian holomorphic bundles, *Special classes of linear operators and other topics (Bucharest, 1986)*, Birkhäuser Verlag, Oper. Theory Adv. Appl., 28, Basel 1988, pp. 267–289.
- [Sz04] R. Szöke, Canonical complex structures associated to connections and complexifications of Lie groups, *Math. Ann.* **329** (2004), no. 3, 553–591.
- [T07] A.B. Tumpach, Hyperkähler structures and infinite-dimensional Grassmannians, *J. Funct. Anal.* **243** (2007), no. 1, 158–206.

- [U85] H. Upmeyer, *Symmetric Banach Manifolds and Jordan C^* -algebras*, North-Holland Mathematics Studies, 104. Notas de Matemàtica, 96. North-Holland Publishing Co., Amsterdam, 1985.
- [W90] D.R. Wilkins, The Grassmann manifold of a C^* -algebra, *Proc. Roy. Irish Acad. Sect. A* **90**(1990), no. 1, 99–116.

INSTITUTE OF MATHEMATICS “SIMION STOILOW” OF THE ROMANIAN ACADEMY, P.O.Box 1-764, RO-014700 BUCHAREST, ROMANIA

E-mail address: `Daniel.Beltita@imar.ro`

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE ZARAGOZA, 50009 ZARAGOZA, SPAIN

E-mail address: `gale@unizar.es`