

A special class of holomorphic mappings and the Faddeev-Hopf model

Radu Slobodeanu

^a Faculty of Physics, Bucharest University, CP Mg-11, RO - 077125.

E-mail: radualexandru.slobodeanu@g.unibuc.ro

Abstract

Pseudo horizontally weakly conformal maps [15] extend both holomorphic and (semi)conformal maps into an almost Hermitian manifold. We find critical points for the (generalized) Faddeev-Hopf model [25] in this larger class.

Keywords: (almost) f -structure, holomorphic map, harmonic map, critical map.

2000 Mathematics Subject Classification: 53C15, 53C43, 53C55, 31B30.

1 Introduction

Harmonic maps between Riemannian manifolds, $\varphi : (M, g) \rightarrow (N, h)$ are critical points for the Dirichlet energy functional (see e.g. [4]):

$$\mathcal{E}(\varphi) = \frac{1}{2} \int_M \|\mathrm{d}\varphi\|^2 v_g.$$

a generalization of the kinetic energy of Classical Mechanics.

Among the most studied classes of harmonic maps are those one that are, in addition, horizontally weakly conformal (HWC) cf. [4] (this class is given precisely by harmonic morphisms). Another well known type of harmonic maps is given by holomorphic maps from a cosymplectic (semi-Kähler) manifold to a (1, 2)-symplectic (quasi-Kähler) one, according to a classical result of Lichnerowicz [14]. Moreover, holomorphic maps between almost Kähler manifolds are minimizers in their homotopy class for the Dirichlet energy.

In this paper we turn our attention to the following energy-type functional proposed in [25] as a generalization of the Faddeev-Hopf static Hamiltonian [11] from hadron physics:

$$\mathcal{E}_{\mathrm{FH}}(\varphi) = \frac{1}{2} \int_M (\|\mathrm{d}\varphi\|^2 + \alpha \|\varphi^* \Omega\|^2) v_g, \quad (1.1)$$

where $\alpha \geq 0$ is a coupling constant and Ω denotes the Kähler 2-form on the target (N, J, h) of the map φ .

The main goal in what follows is to find critical maps for (1.1) that belong to the two classes mentioned above (HWC and holomorphic). As the original model was concerned with the case $M = \mathbb{S}^3$, $N = \mathbb{S}^2$, we are interested to work with a broader notion of holomorphic map that allow also odd dimensional domains. Pseudo horizontally weakly conformal maps (PHWC) into almost Hermitian manifolds, a class that includes also HWC mappings, show to be the appropriate choice.

The paper is organized as follows. The next section survey for further use some known facts about PHWC maps [15], their harmonicity and the corresponding notion of pseudo

horizontally homothetic (PHH) map. Besides we show that a wider perspective is possible by considering mappings from a Riemannian manifold to a f -structured one (rather than just almost complex) and by approaching their harmonicity also from the point of view of conformal geometry. We stress the fact that PHWC property answers the question when we can endow the domain with an almost f -structure compatible with the given metric, such that our map becomes holomorphic in an appropriate sense. Finally, Section 3 investigates the PHWC (stable) critical points for the strong coupling limit of the Faddeev-Hopf model using the variational formulae derived in [25]. Two of the main applications are:

(*) *a holomorphic submersion between Kähler manifolds (or some appropriate analogue) is a critical point for the (full) Faddeev-Hopf functional if it has constant Dirichlet energy along horizontal curves;*

(**) *a Kähler targeted semiconformal (HWC) submersion is a critical point for the strong coupling Faddeev-Hopf functional if and only if it is 4-harmonic.*

2 PHWC mappings

2.1 Generalities about (almost) f -structures

An almost f -structure on M^m is a section F of $\text{End}(TM)$ such that $F^3 + F = 0$. According to [26], the rank k , of F , is even and constant. If $k = m$, then F is an *almost complex structure* on M .

A couple (g, F) is called *metric almost f -structure* if g is a compatible metric, i.e. $g(FX, Y) + g(X, FY) = 0$. In this case we have an *orthogonal* decomposition:

$$T^{\mathbb{C}}M = T^0M \oplus T^{(1,0)}M \oplus T^{(0,1)}M,$$

where T^0M , $T^{(1,0)}M$ and $T^{(0,1)}M$ are the eigenbundles of F corresponding to 0, i and $-i$, respectively. We say that $\mathcal{F} = T^0M \oplus T^{(0,1)}M$ is **the complex distribution associated to F** , cf. [19].

Definition 2.1. ([19]) A map $\varphi : (M, F^M) \rightarrow (N, F^N)$, between manifolds endowed with almost f -structures, is called **holomorphic** if $d\varphi(\mathcal{F}^M) \subseteq \mathcal{F}^N$, where \mathcal{F}^M and \mathcal{F}^N are the complex distributions associated to F^M and F^N , respectively.

In real terms, the above definition says that φ is holomorphic if:

$$d\varphi(F^M X) - F^N d\varphi(X) \in \text{Ker } F^N, \quad \forall X \in TM. \quad (2.1)$$

In particular, when both F^M and F^N are almost complex structures, we find again the well-known definition of a holomorphic map: $d\varphi \circ J^M - J^N \circ d\varphi = 0$.

Definition 2.2. ([19]) An almost f -structure F is called **integrable** if \mathcal{F} is integrable (i.e. for any $X, Y \in \Gamma(\mathcal{F})$ we have $[X, Y] \in \Gamma(\mathcal{F})$); an *f -structure* is an integrable almost f -structure.

Example. If an almost contact structure is *normal* ([5]), then it is integrable in the sense of the above definition, cf. [13] (the converse is not necessarily true). For instance *any* circle bundle over a complex manifold inherits on its total space, a normal almost contact structure.

2.2 The general PHWC condition

The following definition generalizes the corresponding one given in [9, 15] (when the codomain is endowed with an almost complex structure), respectively in [10] (when the codomain is endowed with an almost f -structure):

Definition 2.3. Let (M, g) be a Riemannian manifold and (N, F, h) a manifold endowed with a metric almost f -structure. A map $\varphi : (M, g) \longrightarrow (N, F, h)$ is **pseudo horizontally weakly conformal (PHWC)** if:

$$F \circ [d\varphi \circ d\varphi^t, F] = 0 \quad (2.2)$$

where $d\varphi^t$ stands for the *adjoint map*, $d\varphi_x^t : T_{\varphi(x)}N \rightarrow T_xM$, characterized by: $g(X, d\varphi_x^t(E)) = h(d\varphi_x(X), E)$, $\forall X \in T_xM, E \in T_{\varphi(x)}N$.

The following result generalizes [15, Remark 6]:

Proposition 2.1. *A PHWC map $\varphi : (M, g) \longrightarrow (N, F, h)$ induces a metric almost f -structure, F^φ , on M , with respect to which φ becomes holomorphic.*

Proof. We note firstly that $d\varphi^t(T^{(1,0)}N)$ is a g -isotropic distribution on M .

$$g(d\varphi^t(T^{(1,0)}N), d\varphi^t(T^{(1,0)}N)) = h(T^{(1,0)}N, d\varphi \circ d\varphi^t(T^{(1,0)}N)) = 0,$$

as $d\varphi \circ d\varphi^t(T^{(1,0)}N) \subseteq T^0N \oplus T^{(1,0)}N$, cf. (2.2). According to [23, Prop. 2.2], for any g -isotropic distribution there exists a unique metric f -structure, F^φ , on M , such that $d\varphi^t(T^{(1,0)}N) = \text{Ker}(F^\varphi - i)$. Denote by $T^{\mathbb{C}}M = T^0M \oplus T^{1,0}M \oplus T^{0,1}M$ the corresponding orthogonal splitting. It is easy to remark that: $d\varphi^t(T^{(0,1)}N) = \text{Ker}(F^\varphi + i)$ and $d\varphi(T^0M) \subseteq T^0N$.

It is clear now that φ is holomorphic, according to Definition 2.1, i.e. $d\varphi(T^0M \oplus T^{(0,1)}M) \subseteq T^0N \oplus T^{(0,1)}N$. ■

Remark 2.1. (i) The PHWC condition does not depend on the metric on the codomain but only on the conformal class of the metric on the domain.

(ii) $\text{rank}F^\varphi = \text{rank}F + \text{rank}d\varphi - \dim N$.

In particular, if φ is submersive ($\text{rank}d\varphi = \dim N$), then $\text{rank}F^\varphi = \text{rank}F$. If $F = J$ is an almost complex structure on N ($\text{rank}F = \dim N$), then $\text{rank}F^\varphi = \text{rank}d\varphi$ (we have, moreover, $\text{Ker}F^\varphi = \text{Ker}d\varphi$).

Example. Besides holomorphic maps between almost Hermitian manifolds, the following (already known) classes of mappings satisfy PHWC condition (2.2):

(i) *stable harmonic mappings to an irreducible Hermitian symmetric space of compact type*, cf. [8];

(ii) *(ϕ, J) -holomorphic mappings from a metric almost contact manifold (M, ϕ, ξ, g) to an almost Hermitian manifold (N, J, h)* , i.e. mappings that satisfy $d\varphi \circ \phi = J \circ d\varphi$;

(iii) *semiconformal / horizontally weakly conformal (HWC) mappings*, i.e. mappings that satisfy $d\varphi \circ d\varphi^t = \lambda^2 id$, cf. [4];

(iv) *contact - holomorphic mappings* between almost contact manifolds, cf. [7].

For the sake of simplicity, for the remainder of the article, we will make the assumption that the target is endowed with an almost complex structure. Most of the results in the rest of this section can be easily extended to the general case of PHWC maps to a f -structured manifold.

2.3 PHWC submersions

Suppose now that φ is submersive. The restriction of $d\varphi_x$ to the horizontal space $\mathcal{H}_x = (\text{Ker } d\varphi_x)^\perp$ maps that space isomorphically onto $T_{\varphi(x)}N$. Denote its inverse by $\widehat{\cdot}$; for any $Z \in T_{\varphi(x)}N$, the vector $\widehat{Z} \in \mathcal{H}_x$ is called the *horizontal lift of Z* (this operation can be extended to local vector fields).

As φ takes values in an almost Hermitian manifold (N, J, h) , we have an almost complex structure naturally induced on \mathcal{H} and an almost f -structure, F , that extends it: $F|_{\mathcal{H}}(X) = J_{\mathcal{H}}(X) = J\widehat{d\varphi(X)}$ and $F|_{\mathcal{V}} = 0$, where $\mathcal{V} = \text{Ker } d\varphi$.

In this case, the PHWC condition is equivalent to the compatibility of $J_{\mathcal{H}}$ with the domain metric g (i.e. F is a metric almost f -structure). Indeed, starting with the remark that $d\varphi^t : TN \rightarrow \mathcal{H}$ is this time an isomorphism, we obtain (for an arbitrary compatible metric h on N):

$$\begin{aligned} g(J_{\mathcal{H}}X, d\varphi^t E) &= h(d\varphi(J_{\mathcal{H}}X), E) = h(J(d\varphi X), E) = -h(d\varphi X, JE) \\ &= -g(X, d\varphi^t JE) = -g(X, d\varphi^{-1}(d\varphi(d\varphi^t JE))) \\ &= -g(X, d\varphi^{-1}(J(d\varphi(d\varphi^t E)))) = -g(X, J_{\mathcal{H}}d\varphi^t E). \end{aligned}$$

In this case, an alternative terminology for PHWC is *horizontally holomorphic* cf. [22].

Remark 2.2. (i) If (N, J) is a *complex* manifold (i.e. J is integrable), then the almost f -structure induced by a PHWC submersion $\varphi : (M, g) \rightarrow (N, J)$ is integrable, according to [21, Remark 2.2]. Moreover, *all* f -structures (locally) appear in this way.

(ii) Let $(z^\alpha)_{\alpha=\overline{1, n}}$ be local complex coordinates on the complex manifold (N, J) . The PHWC condition is equivalent to $g^{ij} \frac{\partial \varphi^\alpha}{\partial x^i} \frac{\partial \varphi^\beta}{\partial x^j} = 0, \forall \alpha, \beta$, cf. [15, Lemma 3].

(iii) If the fibers of φ are 1-dim. and M is orientable, we have seen [24] that the induced f -structure is an almost contact metric structure, ϕ . It is integrable but not necessarily normal. The supplementary condition that assures the normality is given in [24, Theorem 4.1].

2.4 Harmonicity of PHWC maps

A mapping is harmonic if and only if its tension field vanishes, cf. [4]. The tension field of a PHWC map into an almost Hermitian manifold (N, J, h) is given by

$$\tau(\varphi) = J\text{div}^\varphi J - d\varphi(F\text{div}F), \quad (2.3)$$

where $\text{div}^\varphi J = \text{trace}_g \varphi^* \nabla^N J$ and $\text{div}F = \text{trace } \nabla F$.

Therefore a PHWC mapping to an (1,2)-symplectic manifold (i.e. $(\nabla_X J)(Y) + (\nabla_{JX} J)(JY) = 0$) will be harmonic if and only if $F\text{div}F = 0$, cf. [17]. If $F\text{div}F = 0$, then I shall call the (almost) f -structure, F , *cosymplectic*.

For submersions, if we consider a local adapted frame $\{e_i, Fe_i, e_\alpha\}$ (i.e. an orthonormal frame such that $e_\alpha \in \text{Ker } F, \forall \alpha$), then the above relation reads:

$$\begin{aligned} \tau(\varphi) &= \sum_{i=1}^n J \left((\nabla_{e_i}^\varphi J)(d\varphi(e_i)) + (\nabla_{Fe_i}^\varphi J)(d\varphi(Fe_i)) \right) \\ &\quad - d\varphi \left(F [(\nabla_{e_i} F)(e_i) + (\nabla_{Fe_i} F)(Fe_i)] + (m - 2n)\mu^\mathcal{V} \right), \end{aligned} \quad (2.4)$$

where $\mu^\mathcal{V}$ denotes the mean curvature of the fibres.

2.5 Pseudo horizontally homothetic maps

Let $\varphi : (M, g) \longrightarrow (N, J, h)$ be a PHWC submersion with minimal fibres onto a (1,2)-symplectic manifold. If the induced f -structure F , on M , satisfies

$$F((\nabla_X F)(X) + (\nabla_{FX} F)(FX)) = 0, \quad \forall X \in \text{Ker}(F^2 + I), \quad (2.5)$$

then φ has the following property: if P is a complex submanifold of N , then $K = \varphi^{-1}(P) \subset M$ is a f -invariant minimal submanifold of M , cf. [17, Prop. 7].

In [3], the same property was proved for *pseudo horizontally homothetic* (PHH) submersions. Recall that a map is PHH if by definition it is PHWC and satisfies $[\text{d}\varphi \circ \nabla_X^M \circ \text{d}\varphi^t, J] = 0, \quad \forall X \in \text{Ker}(F^2 + I)$. For submersions, this means that $J_{\mathcal{H}}$ is $\nabla^{\mathcal{H}}$ -parallel in horizontal directions. Standard examples of PHH mappings are holomorphic mappings from a Sasakian (or Kählerian) domain to a Kähler target.

A PHH submersion with minimal fibres is a (weakly) stable harmonic map, cf. [1]. Further properties and examples of PHH harmonic submersions can be found in [2, 6].

For horizontally weakly conformal maps both conditions (2.5) and PHH reduce to horizontal homothety (HH), i.e. the conformal factor λ is constant in horizontal directions.

2.6 Conformal geometry viewpoint

Let $(M, [g])$ be a conformal manifold ($[g]$ denotes a conformal class of Riemannian metrics).

A **Weyl connection** D on $(M, [g])$ is a torsion-free linear connection which preserves the conformal class $[g]$, cf. [12] (in this case we say that D defines a *Weyl structure* on M). Preserving the conformal class means that for any $\tilde{g} \in [g]$, there exists a 1-form $\theta_{\tilde{g}}$ (called the *Higgs field*) such that:

$$D\tilde{g} = -2\theta_{\tilde{g}} \otimes \tilde{g}.$$

This formula is conformally invariant in the sense that, if $\tilde{g} = e^{2f}g$, then $\theta_{\tilde{g}} = \theta_g - \text{d}f$.

Conversely, if one starts with a fixed Riemannian metric g on M (with ∇^g the Levi-Civita connection) and a fixed 1-form θ (with θ^\sharp the dual vector field with respect to g), then the connection

$$D_X Y = \nabla_X^g Y + \theta(X)Y + \theta(Y)X - g(X, Y)\theta^\sharp, \quad (2.6)$$

is a Weyl connection, preserving the conformal class of g . Clearly, (g, θ) and $(e^{2f}g, \theta - \text{d}f)$ define the same Weyl structure. If, moreover, θ is an exact 1-form, then D is the Levi-Civita connection of some representant of the conformal class.

Suppose now (M^m, g) endowed with a metric almost f -structure. Consider the Weyl structure (2.6) constructed with an arbitrary 1-form on M . Then, with respect to an adapted orthonormal frame $\{e_i, Fe_i, e_\alpha\}$ (i.e. e_α 's span $\text{Ker}F$), we have:

$$\begin{aligned} F \text{div}^D F &= \sum_{i, \alpha} F [(D_{e_i} F)(e_i) + (D_{Fe_i} F)(Fe_i) + (D_{e_\alpha} F)(e_\alpha)] \\ &= F \text{div}^{\nabla^g} F + (m - 2)F^2 \theta^\sharp. \end{aligned}$$

So if I take $\theta^\sharp = \frac{1}{m-2} F \operatorname{div}^{\nabla^g} F$, then D , the Weyl connection associated to the dual 1-form θ , will have the property that:

$$F \operatorname{div}^D F = 0.$$

Notice that D is not uniquely determined. Analogously to [18, Def. 4.2] for the almost complex case, we introduce the following:

Definition 2.4. A Weyl connection on $(M, [g], F)$ will be called **Weyl connection compatible with F** if $F \operatorname{div}^D F = 0$.

Recall that we have an extended notion of harmonicity for maps defined on a conformal manifold [18, Def. 2.1]. Analogously to [18, Prop. 4.5], we have:

Proposition 2.2. A PHWC map $\varphi : (M, g) \rightarrow (N, J, h)$ to a $(1,2)$ -symplectic manifold is harmonic with respect to some compatible Weyl connection on $(M, [g], F^\varphi)$.

3 Critical points for the Faddeev-Hopf functional

The strong coupling limit for the (generalized) *Faddeev-Hopf model* involves the variational problem for the following energy-type functional, cf. [25]:

$$\mathcal{E}_{\text{FH}}^\infty(\varphi) := \lim_{\alpha \rightarrow \infty} \alpha^{-1} \mathcal{E}_{\text{FH}}(\varphi) = \|\varphi^* \Omega\|_{L^2}^2 = \frac{1}{2} \int_M \langle \varphi^* \Omega, \varphi^* \Omega \rangle v_g, \quad (3.1)$$

where $\varphi : (M, g) \rightarrow (N, J, h)$ are mappings defined on a compact, oriented Riemannian manifold, with values in a Kähler manifold with the *fundamental 2-form* $\Omega = h(J\cdot, \cdot)$. According to [25, Corollary 2.4], such a mapping will be a critical point for this functional if and only if:

$$(\delta \varphi^* \Omega)^\sharp \in \operatorname{Ker} d\varphi. \quad (3.2)$$

We can remark that if the target is only a symplectic manifold, the above result is still true (so in the following the Kähler hypothesis can be weakened).

The goal of this section is to identify PHWC submersions with Kähler targets, that satisfy (3.2). Unless otherwise stated, all the mappings considered in the rest of the paper will be Kähler targeted.

Let us begin with an easy to check formula that will be useful in what follows:

Lemma 3.1. Let φ be a submersion between Riemannian manifolds, and let Y, Z be horizontal (local) vector fields on the domain. Then, for any vector field X , we have:

$$(\nabla_X \varphi^* h)(Y, Z) = h(\nabla d\varphi(X, Y), d\varphi(Z)) + h(d\varphi(Y), \nabla d\varphi(X, Z)). \quad (3.3)$$

Proposition 3.1. Let $\varphi : (M, g) \rightarrow (N, J, h)$ be a PHWC submersion from a compact, oriented Riemannian manifold. Then any two of the following statements imply the third:

- (i) The f -structure, F , induced on M is cosymplectic (i.e. $F \operatorname{div} F = 0$)
- (ii) φ is a critical point of $\mathcal{E}_{\text{FH}}^\infty$ (i.e. it verifies the equation (3.2))
- (iii) With respect to any (local) adapted orthonormal frame $\{E_j, FE_j, E_\alpha\}$ on M , the pullback of the codomain metric satisfies:

$$\sum_j [(\nabla_{E_j} \varphi^* h)(FE_j, Z) - (\nabla_{FE_j} \varphi^* h)(E_j, Z)] = 0, \quad \forall Z \in (\operatorname{Ker} d\varphi)^\perp.$$

Proof. The PHWC condition assures us that: $\hat{\Omega}(X, Y) := \varphi^* \Omega(X, Y) = \varphi^* h(FX, Y)$. In order to compute its co-differential, we start with:

$$\begin{aligned}
(\nabla_X \hat{\Omega})(Y, Z) &= X[\varphi^* h(FY, Z)] - \varphi^* h(F \nabla_X Y, Z) - \varphi^* h(FY, \nabla_X Z) \\
&= [h(\nabla_X^\varphi d\varphi(FY), d\varphi(Z)) + h(d\varphi(FY), \nabla_X^\varphi d\varphi(Z))] \\
&\quad - \varphi^* h(F \nabla_X Y, Z) - \varphi^* h(FY, \nabla_X Z) \\
&= \varphi^* h((\nabla_X F)Y, Z) + h(\nabla d\varphi(X, FY), d\varphi(Z)) + h(d\varphi(FY), \nabla d\varphi(X, Z)) \\
&= \varphi^* h((\nabla_X F)Y, Z) + (\nabla_X \varphi^* h)(FY, Z), \quad \forall X, Y, Z \in \text{Ker}(F^2 + I).
\end{aligned}$$

With respect to a (local) adapted orthonormal frame $\{E_j, FE_j, E_\alpha\}$ (i.e. $E_\alpha \in \text{Ker } F$), for any horizontal vector field Z on M , we have:

$$\begin{aligned}
-\delta \hat{\Omega}(Z) &= \sum_{j, \alpha} \iota_{E_j} (\nabla_{E_j} \hat{\Omega})(Z) + \iota_{FE_j} (\nabla_{FE_j} \hat{\Omega})(Z) + \iota_{E_\alpha} (\nabla_{E_\alpha} \hat{\Omega})(Z) \\
&= \varphi^* h(\text{div} F, Z) + \sum_j [(\nabla_{E_j} \varphi^* h)(FE_j, Z) - (\nabla_{FE_j} \varphi^* h)(E_j, Z)].
\end{aligned} \tag{3.4}$$

On the other hand, $\delta \hat{\Omega}(Z) = g((\delta \varphi^* \Omega)^\sharp, Z)$. As φ is a submersion, $\text{Ker } d\varphi = \text{Ker } F$ and our conclusion easily follows. ■

Note that any submersive harmonic PHWC map which satisfies (iii) from Proposition 3.1 will be critical point of the *full* Faddeev-Hopf functional (1.1) (i.e. for every value of the coupling, α , not just the infinite coupling limit).

Corollary 3.1. *A PHWC submersion that satisfies*

$$\nabla d\varphi(X, T^{(0,1)}M) \subseteq \varphi^{-1}T^{(1,0)}N, \quad \forall X \in \Gamma(\mathcal{H}) \tag{3.5}$$

is a critical point for \mathcal{E}_{FH}^∞ if and only if it has minimal fibres.

Proof. The condition (3.5) assures us that $\nabla d\varphi(X, FY) = -J \nabla d\varphi(X, Y)$ which implies immediately the condition (iii) from Proposition 3.1.

On the other hand, as N is endowed with a Kähler structure, from the following easy to check relation (true for all $X, Y \in \Gamma(\mathcal{H})$):

$$0 = (\nabla_X^\varphi J) d\varphi(Y) = d\varphi((\nabla_X F)Y) + \nabla d\varphi(X, FY) - J \nabla d\varphi(X, Y), \tag{3.6}$$

we can deduce that $d\varphi((\nabla_X F)Y) + 2\nabla d\varphi(X, FY) = 0$. In particular, we also have: $d\varphi((\nabla_X F)X + (\nabla_{FX} F)FX) = 0$ and therefore the condition (i) from Proposition 3.1 is also satisfied if and only if φ has minimal fibres. ■

Notice that a map as above will satisfy (2.5) and will be (1, 1)-geodesic (see [4]).

As the strong coupling term of the Faddeev model comprises fourth power derivative terms, we expect some resemblance to the case of 4-energy. Let us recall that *p-harmonic maps* are critical points for the *p*-energy $\mathcal{E}_p(\varphi) = \frac{1}{p} \int_M \|d\varphi\|^p v_g$ (see e.g. [16] and references therein).

Corollary 3.2. (i) A PHH submersion is a critical point for the Faddeev-Hopf functional \mathcal{E}_{FH}^∞ if and only if $\operatorname{div}\varphi^*h|_{\mathcal{H}} = 0$.

(ii) A PHH submersion with $\operatorname{grad}^{\mathcal{H}}\|\mathrm{d}\varphi\|^2 = 0$ is a critical point of the Faddeev-Hopf functional \mathcal{E}_{FH}^∞ if and only if it has minimal fibres. In this case it is moreover harmonic and 4-harmonic.

Proof. By a similar computation as for the proof of Proposition 3.1, using (3.6) we can check that:

$$(\nabla_X \hat{\Omega})(Y, Z) = -(\nabla_X \varphi^*h)(Y, FZ) - \varphi^*h(Y, (\nabla_X F)Z), \quad \forall X, Y, Z \in \operatorname{Ker}(F^2 + I).$$

Therefore:

$$\delta \hat{\Omega}(Z) = \operatorname{div}\varphi^*h(FZ) + \sum_j [\varphi^*h(E_j, (\nabla_{E_j} F)Z) + \varphi^*h(FE_j, (\nabla_{FE_j} F)Z)].$$

As our PHH hypothesis (i.e. F parallel in horizontal directions) assures the cancellation of every term in the above sum, the conclusion follows.

For the second statement, use the fact that a PHH submersion is harmonic if and only if it has minimal fibres and then take into account that a submersion is harmonic if and only if its stress-energy tensor is divergence free: $\operatorname{div}S_\varphi = (1/2)\|\mathrm{d}\varphi\|^2 - \operatorname{div}\varphi^*h = 0$. ■

Example. (i.) A holomorphic submersion from a Kähler manifold (or a compact Vaisman manifold) onto a Kähler manifold, which has constant energy density (in horizontal directions), is a critical point of the full Faddeev-Hopf functional (1.1).

(ii.) A Boothby-Wang fibration of a compact, regular Sasakian (or just K-contact) manifold over a Kähler (or just almost Kähler) manifold is a critical point of the full Faddeev-Hopf functional (1.1) (on the total space we consider the metric $g = \varphi^*h + \eta \otimes \eta$ induced from the base, see [5]).

For semiconformal (HWC) particular case, a more precise statement can be made, providing us with class of examples that extend [25, Examples 3.1, 3.3]:

Proposition 3.2. A semiconformal (HWC) submersion $\varphi : (M^m, g) \rightarrow (N^{2n}, J, h)$ with dilation λ is a critical point for the Faddeev-Hopf functional \mathcal{E}_{FH}^∞ if and only if:

$$(2n - 4)\operatorname{grad}^{\mathcal{H}}(\ln\lambda) + (m - 2n)\mu^\mathcal{V} = 0. \quad (3.7)$$

that is, if and only if it is 4-harmonic (morphism, [16]).

Proof. Semiconformal submersions are in particular PHWC, so we always have an induced metric almost f -structure, F , on M and, in addition:

$$F \operatorname{div}F = (2n - 2)\operatorname{grad}^{\mathcal{H}}(\ln\lambda) + (m - 2n)\mu^\mathcal{V}. \quad (3.8)$$

As $\varphi^*h|_{\mathcal{H} \times \mathcal{H}} = \lambda^2 g|_{\mathcal{H} \times \mathcal{H}}$, in this case we shall have, for any Y, Z horizontal vector fields :

$$(\nabla_X \hat{\Omega})(Y, Z) = X(\lambda^2)g(FY, Z) + \lambda^2 g((\nabla_X F)Y, Z).$$

Then, taking the trace w.r.t. an adapted orthonormal frame, we obtain:

$$-\delta \hat{\Omega}(Z) = \lambda^2 g(F \operatorname{div}F - 2\operatorname{grad}^{\mathcal{H}}(\ln\lambda), FZ) \quad (3.9)$$

Now taking into account also (3.8), the conclusion follows. ■

Corollary 3.3. (i). *A horizontally homothetic submersion is a stationary point for the full Faddeev-Hopf functional (1.1) if and only if it has minimal fibres.*

(ii). *A semiconformal submersion onto a four-manifold is a stationary point for the full Faddeev-Hopf functional (1.1) if and only if it has minimal fibres.*

3.1 Stability

According to [25, Corollary 4.9], the Hessian of a critical φ for the energy $\mathcal{E}_{\text{FH}}^\infty$ is

$$\text{Hess}_\varphi(v, v) = \|\varphi^* d(\iota_v \Omega)\|_{L^2}^2 + \int_M \Omega(v, \nabla_{Z_\varphi}^\varphi v) v_g, \quad \forall v \in \Gamma(\varphi^{-1}TN)$$

where $Z_\varphi = (\delta\varphi^*\Omega)^\sharp$.

Therefore a sufficient condition for a critical point of $\mathcal{E}_{\text{FH}}^\infty$ to be (weakly) stable is the second term of $\text{Hess}_\varphi(v, v)$ to be nonnegative. In particular, if $\delta\varphi^*\Omega(V) = 0, \forall V \in \text{Ker } d\varphi$, this second term will vanish. But for a PHWC submersion we can check that $(\nabla_X \varphi^*\Omega)(Y, V) = -\varphi^*h(FY, \nabla_X V)$ and therefore

$$-\delta\varphi^*\Omega(V) = \sum_{i=1}^n \lambda_i^2 g([E_i, FE_i], V), \quad \forall V \in \text{Ker } d\varphi,$$

where λ_i^2 are the nonzero eigenvalues of φ^*h with respect to g and $\{E_i, FE_i\}$ a frame of horizontal eigenvector fields (which exists around almost any point of M , cf. [22]). We can easily conclude that a critical PHWC submersion for $\mathcal{E}_{\text{FH}}^\infty$ is (weakly) stable if any of the following statements holds good:

- (a) the horizontal distribution \mathcal{H} is integrable;
- (b) the induced f -structure on M satisfies

$$(\nabla_X F)(X) + (\nabla_{FX} F)(FX) = 0, \quad \forall X \in \Gamma(\mathcal{H}).$$

On the other hand, as basic vector fields locally span the horizontal distribution of a submersion φ , for any $v \in \Gamma(\varphi^{-1}TN)$, exists X_v local horizontal vector field on M such that $v = d\varphi(X_v)$. In this case, as $\nabla_V^\varphi d\varphi(X) = d\varphi([V, X])$ we have

$$\Omega(v, \nabla_V^\varphi v) = \varphi^*h(FX_v, [V, X_v]), \quad \forall V \in \text{Ker } d\varphi. \quad (3.10)$$

If this term vanishes our submersive critical point will be stable.

Let us illustrate these situations with the following result (for the notions related to contact geometry we refer to [5]).

Proposition 3.3. *Let $\varphi : (M^{2n+1}, \phi, \xi, \eta, g) \rightarrow (N^{2n}, J, h)$ be a (ϕ, J) -holomorphic, horizontally homothetic submersion from an almost contact metric manifold to a (almost) Kähler one. Then φ is a stable critical point for $\mathcal{E}_{\text{FH}}^\infty$ in any of the following cases:*

- (i) (ϕ, ξ, η, g) is a nearly cosymplectic structure;
- (ii) (ϕ, ξ, η, g) is a Kenmotsu structure;
- (iii) (ϕ, ξ, η, g) is a Sasakian structure.

Proof. For horizontally homothetic submersions Proposition 3.1 - (iii) always holds. In all three cases ϕ is, in particular, a cosymplectic f -structure, so we can apply Proposition 3.1 (for $\phi = F$) and conclude that the map is critical.

Nearly cosymplectic structures are defined by $(\nabla_X \phi)X = 0$, so the above (b) condition is true, providing stability. For Kenmotsu manifolds, the contact distribution (which coincides with \mathcal{H}) is integrable, so the above (a) condition is true.

In the last case, as $\mathcal{L}_\xi d\eta = 0$ and $g(X, \phi Y) = d\eta(X, Y)$, we can use (3.10) to show that the second term of the Hessian is equal to zero. ■

Notice that the almost contact structure on M could be seen as the induced f -structure by φ , as a horizontally homothetic map is automatically PHWC (in fact it is moreover PHH, as the target is supposed to be Kähler).

Corollary 3.4. *Boothby-Wang fibrations considered in Example 3-(ii) (in particular the Hopf fibration) are stable critical points for \mathcal{E}_{FH}^∞ .*

Acknowledgments. *I am grateful to Radu Pantilie and Liviu Ornea for very useful remarks and encouraging discussions. I acknowledge the support by the CEx grant no. 2-CEx 06-11-22/25.07.2006.*

References

- [1] Aprodu M. A., PHH harmonic submersions are stable, math.DG/0504069.
- [2] Aprodu M. A. and Aprodu M., Implicitly defined harmonic PHH submersions, *Manuscripta Math.* **100** (1999), 103–121.
- [3] Aprodu M. A., Aprodu M. and Brînzănescu V., A class of harmonic submersions and minimal submanifolds, *Int. J. Math.* **11**(9) (2000), 1177–1191.
- [4] Baird P. and Wood J.C., *Harmonic Morphisms Between Riemannian Manifolds*, Clarendon Press - Oxford, 2003.
- [5] Blair D.E., *Riemannian Geometry of Contact and Symplectic Manifolds*, Birkhauser Boston, Progress in Mathematics, vol.203, 2002.
- [6] Brînzănescu V., Pseudo-harmonic morphisms; applications and examples, *An. Univ. Timișoara Ser. Mat.-Inform.* **39** (2001), Special Issue: Mathematics, 111–121.
- [7] Brînzănescu V. and Slobodeanu R., Holomorphicity and Walczak formula on Sasakian manifolds, *J. Geom. Phys.* **57** (2006), 193–207.
- [8] Burns D., Burstall F., De Bartolomeis P. and Rawnsley J., Stability of harmonic maps of Kähler manifolds, *J. Diff. Geom.* **30** (1989), 579–594.
- [9] Chen J., Structures of certain harmonic maps into Kähler manifolds, *Int. J. Math.* **8** (1997), 573–581.

-
- [10] Erdem S., φ -pseudo harmonic morphisms, some subclasses and their liftings to tangent bundles, *Houston J. Math.* **30** (2004), 1009–1038.
- [11] Faddeev L. D. and Niemi A. J., Stable knot-like structures in classical field theory, *Nature* **387** (1997), 58–61.
- [12] Gauduchon P., Structures de Weyl et théorèmes d’annulation sur une variété conforme autoduale, *Ann. Sc. Norm. Sup. Pisa* **XVIII** **4** (1991), 563–629.
- [13] Ianoş S., Sulle varietà di Cauchy-Riemann, *Rend. Dell’Accademia de Scienze di Napoli*, **39**(1972), 191–195.
- [14] Lichnerowicz A., Applications Harmoniques et Variétés Kähleriennes, *Sympos. Math.* **III** (1970), 341–402.
- [15] Loubeau E., Pseudo harmonic morphisms, *Int. J. of Math.* **8** (1997), 943–957.
- [16] Loubeau E., On p -harmonic morphisms, *Diff. Geom. and Appl.* **12** (2000), 219–229.
- [17] Loubeau E. and Mo X., The geometry of pseudo harmonic morphisms, *Beiträge Algebra Geom.* **45**(1) (2004), 87–102.
- [18] Loubeau E. and Pantilie R., Harmonic morphisms between Weyl spaces and twistorial maps, *Comm. Anal. Geom.* **14** (2006), 847–881.
- [19] Loubeau E. and Pantilie R., Harmonic morphisms between Weyl spaces and twistorial maps II, math.DG/0610676.
- [20] Pantilie R., On submersive harmonic morphisms, in *Harmonic morphisms, harmonic maps, and related topics* (Brest, 1997), Chapman & Hall/CRC Res. Notes Math., **413**, 2000, 23–29.
- [21] Pantilie R., On a class of twistorial maps, *Diff. Geom. Appl.* **26** (2008), 366–376.
- [22] Pantilie R. and Wood J.C., Harmonic morphisms with one-dimensional fibres on Einstein manifolds, *Trans. Amer. Math. Soc.* **354** (2002), 4229–4243.
- [23] Rawnsley J. H., f -structures, f -twistor spaces and harmonic maps, in *Geometry seminar "Luigi Bianchi" II - 1984*, , Lecture Notes in Math., **1164**, Springer, Berlin, 1985, 85–159.
- [24] Slobodeanu R., Pseudo-harmonic morphisms with low dimensional fibers, *Rend. Circ. Mat. Palermo*, **II(LV)** (2006), 5–20.
- [25] Speight J. M., Svensson M., On the Strong Coupling Limit of the Faddeev-Hopf Model, *Commun. Math. Phys.* **272** (2007), 751–773.
- [26] Stong R. E. , The rank of an f -structure, *Kodai Math. Sem. Rep.* **29** (1977), 207–209.